

## Datasheet: PBP015KZZ

Description:	BOVINE DENDRITIC CELL GROWTH KIT
Name:	BOVINE DENDRITIC CELL GROWTH KIT
Format:	Kit
<b>Product Type:</b>	Kits
Quantity:	5 ml

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Functional Assays	-			1:20

Where this reagent has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the protein for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Product Form	Mixed recombinant bovine Interleukin-4 and bovine GM-CSF – supplied as a liquid
Preparation	Recombinant cytokines expressed in mammalian Chinese Hamster Ovary (CHO) cells using the pEE14® vector grown in antibiotic free media and USDA-approved dialysed FCS which has beer screened for BVDV and virus growth by PCR.
Preservative Stabilisers	None present
Endotoxin Level	<0.5EU/mL

#### **Product Information**

**Bovine dendritic cell growth kit** contains a cocktail of biologically active interleukin-4 (IL-4) and granulocyte/macrophage-colony stimulating factor (GM-CSF) that have been premixed at optimal concentrations to induce dendritic cell development from peripheral blood-derived bovine (cattle) monocytes.

#### References

- 1. Hope, J.C. *et al.* (2000) Dendritic cells induce CD4+ and CD8+ T-cell responses to *Mycobacterium bovis* and *M. avium* antigens in *Bacille Calmette Guérin* vaccinated and nonvaccinated cattle. <u>Scand J Immunol. 52 (3): 285-91.</u>
- 2. Walters, A.A. *et al.* (2015) Assessment of the enhancement of PLGA nanoparticle uptake by dendritic cells through the addition of natural receptor ligands and monoclonal antibody. <u>Vaccine.</u> pii: S0264-410X(15)01549-2.

### **Further Reading**

1. Werling, D. *et al.* (1999) Involvement of caveolae in the uptake of respiratory syncytial virus antigen by dendritic cells. <u>J Leukoc Biol. 66 (1): 50-8.</u>

#### Recommended Protocol

- 1. Prepare peripheral blood mononuclear cells (PBMC) from heparinised blood by density gradient centrifugation.
- 2. Purify CD14<sup>+ve</sup> cells by labelling PBMC with CD14 mAb and utilise magnetic bead or flow cytometric separation techniques.
- 3. Resuspend the isolated CD14 $^{+ve}$  cells at a concentration of 1x10 $^{6}$  cells/ml in tissue culture medium (TCM = RPMI or equivalent + 10% foetal calf serum) containing a final dilution of 1:20 of PBP015KZZ .
- 4. Add 3ml of cell suspension to each well of a 6 well tissue culture plate.
- 5. Culture cells in a humidified atmosphere of 5% CO<sub>2</sub> in air, at approximately 37°C.
- 6. Culture cells for 3 days. The cells may then be harvested and used for other procedures including immunophenotyping (as required).
- 7. If a longer culture period is required the cells must be 'fed' with new TCM containing cytokines on day 3:

Carefully remove 1ml spent medium from each well, care is required to avoid disturbing the cells. Add 1.5ml fresh, pre-warmed TCM containing cytokines at 1:20 to each well and re-culture the DC for required culture period (typically up to 7 days).

8. At the end of the culture period adherent and non-adherent cells can be pooled for use in immunoassays and phenotyped (as required). Adherent cells may require a dissociation step to remove them from the plate.

#### Storage

Store at -20°C only.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature this recombinant protein. Should this product contain a precipitate we recommend microcentrifugation before use.

#### Shelf Life

6 months from date of despatch.

## Acknowledgements

This reagent was produced as part of the BBSRC/SEERAD Immunological Toolbox. The kit development was also supported by the European Community's Seventh Framework Programme (FP7, 2007-2013), Research Infrastructures action, under the grant agreement No. FP7-228394 (NADIR)

# Health And Safety Information

Material Safety Datasheet documentation #10286 available at: 10286: https://www.bio-rad-antibodies.com/uploads/MSDS/10286.pdf

## Regulatory

For research purposes only

## Related Products

#### **Recommended Useful Reagents**

MOUSE ANTI BOVINE CD14:FITC (MCA2678F)

MOUSE ANTI HUMAN CD14:Low Endotoxin (MCA1568EL)

MOUSE ANTI HUMAN CD14:Alexa Fluor® 647 (MCA1568A647)

MOUSE ANTI HUMAN CD14:Biotin (MCA1568B)

MOUSE ANTI HUMAN CD14:FITC (MCA1568F)

MOUSE ANTI HUMAN CD14:Pacific Blue® (MCA1568PB)

MOUSE ANTI HUMAN CD14:RPE (MCA1568PE)

MOUSE ANTI HUMAN CD14:Alexa Fluor® 700 (MCA1568A700)

MOUSE ANTI HUMAN CD14:RPE-Alexa Fluor® 647 (MCA1568P647)

MOUSE ANTI BOVINE MHC CLASS II DQ (MCA5655)

MOUSE ANTI BOVINE MHC CLASS II DR (MCA5656)

MOUSE ANTI BOVINE MHC CLASS II DQ:FITC (MCA5655F)

MOUSE ANTI BOVINE CD1w2 (MCA831G)

North & South Tel: +1 800 265 7376

America

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

'M316693:180607'

#### Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint