

## Datasheet: 102002

Description:	GOAT ANTI MOUSE IgM:FITC
Specificity:	IgM
Format:	FITC
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	1 mg

## **Product Details**

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution Flow Cytometry . 1/100 Immunohistology - Frozen Immunohistology - Paraffin . ELISA . Western Blotting Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. **Target Species** Mouse **Product Form** Purified Ig conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid Max Ex/Em Fluorophore Excitation Max (nm) **Emission Max (nm)** FITC 490 525 Antiserum Preparation Antiserum to mouse IgM was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography on mouse IgM covalently linked to agarose. **Buffer Solution** Phosphate buffered saline Preservative 0.1% Sodium azide **Stabilisers** Approx. Protein Ig concentration 1.0 mg/ml Concentrations Immunogen Mouse IgM paraproteins

External Database	UniProt:
Links	P01872 Related reagents
	P01873 Related reagents
	Entrez Gene: <u>16019</u> Ighm <u>Related reagents</u>
	<u>16019</u> Ighm <u>Related reagents</u>
Specificity	<b>Goat anti mouse IgM antibody</b> recognises the heavy chain of mouse IgM as demonstrated by ELISA and flow cytometry. Minimal cross reactivity is observed with human immunoglobulins.
	Goat anti mouse IgM antibody has been cross absorbed against Mouse $IgG_1$ , $IgG_{2a}$ , $IgG_{2b}$ , $IgG$ and $IgA$ , pooled human sera and purified human paraproteins.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
References	1. Kerr, K. <i>et al.</i> (2010) Inflammatory cytokine responses in a pregnant mouse model of <i>Chlamydonbila abortus</i> infection. Vet Microbiol. 144 (3.4): 392.8
	Chlamydophila abortus infection. <u>Vet Microbiol. 144 (3-4): 392-8.</u> 2. Ferrian, S. <i>et al.</i> (2012) Effect of high temperature on blood lymphocyte populations in two
	different genetic rabbit lines. <u>Proceedings 10 th World Rabbit Congress 1169-73</u>
	3. Kamat, M.M. et al. (2016) Changes in Myeloid Lineage Cells in the Uterus and Peripheral Blo
	o. Ramat, M.M. et al. (2010) Onanges in Mycloid Eineage Oens in the Oterus and Perpheral Die
	of Dairy Heifers During Early Pregnancy. Biol Reprod. 95 (3): 68.
	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell</u>
	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell</u> <u>Res. 355 (1): 26-39.</u>
	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Sterr</li> </ul>
	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell</u> <u>Res. 355 (1): 26-39.</u>
	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> </ul>
Storage	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> <li>Store at +4°C or at -20°C if preferred.</li> </ul>
Storage	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Sterr Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> <li>Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended.</li> </ul>
Storage	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19 (2)Feb 10 [Epub ahead of print].</u></li> <li>Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denaited the stored of the stored undiluted.</li> </ul>
Storage	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Sterr Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> <li>Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended.</li> </ul>
Storage Shelf Life	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> <li>Store at +4°C or at -20°C if preferred.</li> <li>Storage in frost-free freezers is not recommended.</li> <li>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denar the antibody. Should this product contain a precipitate we recommend microcentrifugation before</li> </ul>
	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> <li>Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denait the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</li> <li>Please see label for expiry date.</li> </ul>
Shelf Life Health And Safety	<ul> <li>of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u></li> <li>4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u></li> <li>5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].</li> <li>Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denar the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</li> <li>Please see label for expiry date.</li> </ul>
Shelf Life Health And Safety Information	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell</u> <u>Res. 355 (1): 26-39.</u> 5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print]. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denait the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Please see label for expiry date. Material Safety Datasheet documentation #10303 available at: 10303: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf</u> For research purposes only
Shelf Life Health And Safety Information Regulatory th & South Tel: +1 800 erica Fax: +1 915	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell</u> <u>Res. 355 (1): 26-39.</u> 5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print]. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may dena the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Please see label for expiry date. Material Safety Datasheet documentation #10303 available at: 10303: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf</u> For research purposes only 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 Fax: +44 (0)1865 852 739 Europe Tel: +49 (0) 89 8090 95 21 Fax: +44 (0)1865 852 739 Europe Tel: +49 (0) 89 8090 95 50
Shelf Life Health And Safety Information Regulatory th & South Tel: +1 800 erica Fax: +1 915	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell Res. 355 (1): 26-39.</u> 5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print].         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.         This product should be stored undiluted. Avoid repeated freezing and thawing as this may denait the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.         Please see label for expiry date.         Material Safety Datasheet documentation #10303 available at: 10303: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf</u> For research purposes only         265 7376       Worldwide       Tel: +44 (0)1865 852 700       Europe       Tel: +49 (0) 89 8090 95 21
Shelf Life Health And Safety Information Regulatory th & South Tel: +1 800 erica Fax: +1 915	of Dairy Heifers During Early Pregnancy. <u>Biol Reprod. 95 (3): 68.</u> 4. Monzo, H.J. <i>et al.</i> (2017) Insulin promotes cell migration by regulating PSA-NCAM. <u>Exp Cell</u> <u>Res. 355 (1): 26-39.</u> 5. Desancé, M. <i>et al.</i> (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. <u>Int J Mol Sci. 19</u> (2)Feb 10 [Epub ahead of print]. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may dena the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Please see label for expiry date. Material Safety Datasheet documentation #10303 available at: 10303: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf</u> For research purposes only 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 Fax: +44 (0)1865 852 739 Europe Tel: +49 (0) 89 8090 95 21 Fax: +44 (0)1865 852 739 Europe Tel: +49 (0) 89 8090 95 50

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint