

## Datasheet: 103009

<b>Description:</b>	GOAT ANTI MOUSE IgG:RPE
<b>Specificity:</b>	IgG
<b>Format:</b>	RPE
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.5 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
ELISA			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified Ig conjugated to R. Phycoerythrin (RPE) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>RPE 488nm laser</td> <td>496</td> <td>578</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	RPE 488nm laser	496	578
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
RPE 488nm laser	496	578					
<b>Preparation</b>	Purified Ig prepared by affinity chromatography on pooled mouse IgG covalently linked to agarose						
<b>Antiserum Preparation</b>	Antisera to mouse IgG were raised by repeated immunisations of goats with highly purified antigen.						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative</b>	0.1% Sodium Azide (NaN <sub>3</sub> )						
<b>Stabilisers</b>	Stabilizing agent						
<b>Approx. Protein Concentrations</b>	Ig concentration 0.5 mg/ml						
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P01867</a> <a href="#">Related reagents</a> <a href="#">P01865</a> <a href="#">Related reagents</a> <a href="#">P01863</a> <a href="#">Related reagents</a>						

[P01864](#) [Related reagents](#)  
[P01868](#) [Related reagents](#)  
[P01869](#) [Related reagents](#)  
[P03987](#) [Related reagents](#)

**Entrez Gene:**

[16016](#) Ighg2b [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[16017](#) Ighg1 [Related reagents](#)  
[16017](#) Ighg1 [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[380795](#) AI324046 [Related reagents](#)

---

**Synonyms**

Igh-4

---

**Specificity**

**Goat anti Mouse IgG antibody** recognizes mouse IgG, recognising the heavy chain of mouse IgG1, IgG2a, IgG2b and IgG3 as demonstrated by ELISA.

Goat anti Mouse IgG antibody has been cross-adsorbed against mouse IgM, mouse IgA and human serum to reduce potential cross-reactivity.

---

**References**

1. Joimel, U. *et al.* (2010) Stimulation of angiogenesis resulting from cooperation between macrophages and MDA-MB-231 breast cancer cells: proposed molecular mechanism and effect of tetrathiomolybdate. [BMC Cancer. 10: 375.](#)
2. Childs K *et al.* (2012) Paramyxovirus V proteins interact with the RNA Helicase LGP2 to inhibit RIG-I-dependent interferon induction. [J Virol. 86 \(7\): 3411-21.](#)
3. Moalli, F. *et al.* (2015) Intravital and whole-organ imaging reveals capture of melanoma-derived antigen by lymph node subcapsular macrophages leading to widespread deposition on follicular dendritic cells. [Front Immunol. 6: 114.](#)
4. Ramos-Sevillano, E. *et al.* (2016) PSGL-1 on Leukocytes is a Critical Component of the Host Immune Response against Invasive Pneumococcal Disease. [PLoS Pathog. 12 \(3\): e1005500.](#)
5. Abbate, F. *et al.* (2016) Acid-sensing ion channel immunoreactivities in the cephalic neuromasts of adult zebrafish. [Ann Anat. 207: 27-31.](#)

---

**Storage**

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Shelf Life**

12 months from date of despatch.

---

**Health And Safety Information**

Material Safety Datasheet documentation #10331 available at:  
10331: <https://www.bio-rad-antibodies.com/uploads/MSDS/10331.pdf>

---

**Regulatory**

For research purposes only

---

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751  
Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide** Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739  
Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe** Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50  
Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M314450:180412'

