

## Datasheet: 204002

Description:	GOAT ANTI HUMAN IgG:FITC		
Specificity:	lgG		
Format:	FITC		
Product Type:	Polyclonal Antibody		
lsotype:	Polyclonal IgG		
Quantity:	1 mg		

# Product Details

Applications		This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from						
	the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
	Flow Cytometry	-						
	Immunohistology - Froz	zen 🔹						
	Immunofluorescence	-						
	Where this product has not been tested for use in a particular technique this does not necessarily							
	exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is							
	recommended that the user titrates the product for use in their own system using appropriate							
	negative/positive cor			, , , , , , , , , , , , , , , , , , ,	<b>3 () ()</b>			
Target Species	Human							
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid							
Max Ex/Em	Fluorophore	Excitation Max (n	m) Emis	sion Max (nm)				
	FITC	490		525				
Preparation	Purified IgG prepared by affinity chromatography							
Buffer Solution	Phosphate buffered							
Preservative Stabilisers	0.1% Sodium azide							
Approx. Protein Concentrations	IgG concentration 1.							
Immunogen	Normal human IgG and human IgG paraproteins							
External Database	UniProt							
Links	UniProt:							
	<u>P01857</u> <u>Rela</u>	ted reagents						
	P01859 Rela	ted reagents						
	P01861 Rela	ted reagents						

	P01860 Related reagents					
	Entrez Gene:					
	3500 IGHG1 Related reagents					
	3501 IGHG2 Related reagents					
	3502 IGHG3 Related reagents					
	3503 IGHG4 Related reagents					
Specificity	Goat anti Human IgG antibody reacts with the heavy chain of human IgG and has been cross- absorbed with human IgM, IgD and IgA, but may recognise IgG from other species.					
References	<ol> <li>Lund K <i>et al.</i> (2014) The novel EpCAM-targeting monoclonal antibody 3-17I linked to saporin is highly cytotoxic after photochemical internalization in breast, pancreas and colon cancer cell lines. <u>MAbs. 6 (4): 1038-50.</u></li> <li>Gilbert, A. (2013) Humoral immune response to melanoma: discovery and evaluation of anti-melanoma antibodies <u>PhD. Thesis KCL UK.</u></li> </ol>					
Storage	Store at +4°C or at -20°C if preferred.					
	Storage in frost-free freezers is not recommended.					
	This product should be stored undiluted. This product is photosensitive and should be protected					
	from light.					
	Avoid repeated freezing and thawing as this may denature the antibody.					
Shelf Life	Please see label for expiry date.					
Health And Safety Information	Material Safety Datasheet documentation #10303 available at: 10303: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf</u>					
Regulatory	For research purposes only					

## **Related Products**

#### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

'M314294:180412'

#### Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint