

Datasheet: 2222-6604F

Description:	SHEEP ANTI HUMAN C3c:FITC
Specificity:	C3c
Other names:	COMPLEMENT COMPONENT 3c
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	•			1/500 - 1/50K
Immunofluorescence				1/10 - 1/50

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Human				
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Preparation	Purified IgG prepared by affinity chromatography				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)				
Approx. Protein Concentrations	IgG concentration 1.0	Omg/ml			
Immunogen	Native human C3c purified from serum				
External Database Links	UniProt:				

Related reagents

Entrez Gene:

P01024

718 C3 Related reagents

Synonyms	CPAMD1
Specificity	Sheep anti Human C3c antibody detects human complement 3c, a polypeptide fragment which is present in C3. C3 plays a central role in the activation of the complement system. Processing of C3 by C3 convertase is the central reaction in both classical and alternative complement pathways, resulting in C3a and C3b. C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates, and help to initiate the complement cascade, potentially resulting in cellular apoptosis. C3a anaphylatoxin is a mediator of local inflammatory process. It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes.
	C3b is rapidly split in two positions by factor I and a cofactor to form iC3b (inactivated C3b) and C3f which is released. Then iC3b is slowly cleaved (possibly by factor I) to form C3c and C3dg. Defects in C3 can result in susceptibility to pyogenic infection.
	This antbody shows minimal reactivity with related serum proteins.
References	1. Janssen, B.J. <i>et al.</i> (2005) Structures of complement component C3 provide insights into the function and evolution of immunity. <u>Nature</u> . <u>437 (7058)</u> : <u>505-11</u> .
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M301221:170109'

Printed on 10 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com