

Datasheet: 642009

Description:	DONKEY ANTI GOAT IgG (H/L):RPE (RAT/MOUSE ADSORBED)
Specificity:	IgG (H/L)
Format:	RPE
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Goat		
Species Cross Reactivity	Does not react with: Mouse, Rat		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578

Antiserum Preparation Antiserum to goat IgG (H/L) was raised by repeated immunisation of donkeys with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative	0.1% Sodium azide
Stabilisers	Stabilising agent
Approx. Protein Concentrations	0.5 mg/ml
Immunogen	Goat IgG.

Specificity **Donkey anti Goat IgG antibody** recognizes the heavy and light chains of goat IgG as demonstrated by ELISA. The preparation has minimal cross reaction with rat and mouse serum proteins, however, it may react with the light chains of other goat immunoglobulins.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Gerry, A.B. and Leake, D.S. (2014) Effect of low extracellular pH on NF- κ B activation in macrophages. [Atherosclerosis. 233: 537-44.](#)
2. Armour, K.L.*et al.* (2014) Clearance of Human IgG1-Sensitised Red Blood Cells In Vivo in Humans Relates to the In Vitro Properties of Antibodies from Alternative Cell Lines. [PLoS One. 9: e109463.](#)

Storage +4°C DO NOT FREEZE
This product is photosensitive and should be protected from light.

Shelf Life 18 months from the date of despatch.

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M314456:180412'

Printed on 19 Apr 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)