

## Datasheet: AAI29P

| Description:  | GOAT ANTI CHICKEN IgG (Fc):HRP |
|---------------|--------------------------------|
| Specificity:  | lgG (Fc)                       |
| Other names:  | ΙgΥ                            |
| Format:       | HRP                            |
| Product Type: | Polyclonal Antibody            |
| Isotype:      | Polyclonal IgG                 |
| Quantity:     | 1 mg                           |
|               |                                |

# **Product Details**

| Applications                      |   | This product has been reported to work in the following applications. This information is derived |                           |  |                             |  |  |  |
|-----------------------------------|---|---|---------------------------|--|-----------------------------|--|--|--|
|                                   | from testing within our laboratories, peer-reviewed publications or personal communications from<br>the originators. Please refer to references indicated for further information. For general protocol |   |                           |  |                             |  |  |  |
|                                   | recommendations, please visit www.bio-rad-antibodies.com/protocols.   |   |                           |  |                             |  |  |  |
|                                   | recommendations, please   | Yes   | No                        | Not Determined                                       | Suggested Dilution          |  |  |  |
|                                   | Immunohistology - Frozen  |   |                           | •  |                             |  |  |  |
|                                   | Immunohistology - Paraffin  |   |                           | •  |                             |  |  |  |
|                                   | ELISA   | •   |                           |  | 1/10,000 - 1/100,000        |  |  |  |
|                                   | Western Blotting  |   |                           |  |                             |  |  |  |
|                                   | Where this antibody has not been tested for use in a particular technique this does not ne  |   |                           |  |                             |  |  |  |
|                                   | exclude its use in such procedures. Suggested working dilutions are given as a guide only   |   |                           |  |                             |  |  |  |
|                                   | recommended that the use  |   |                           |  |                             |  |  |  |
|                                   | negative/positive controls.   |   |                           |  |                             |  |  |  |
| Target Species                    | Chicken   |   |                           |  |                             |  |  |  |
| Product Form                      | Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid  |   |                           |  |                             |  |  |  |
| Antiserum Preparatio              | <b>on</b> Antisera to chicken IgG we<br>Purified IgG prepared by a  |   | • •                       | •  | th highly purified antigen. |  |  |  |
| Buffer Solution                   | Phosphate buffered saline   |   |                           |  |                             |  |  |  |
| Preservative                      | 0.05% Proclin™ 300  |   |                           |  |                             |  |  |  |
| Stabilisers                       | 0.2% Bovine Serum Albumin   |   |                           |  |                             |  |  |  |
| Approx. Protein<br>Concentrations | IgG concentration 1.0 mg/ml   |   |                           |  |                             |  |  |  |
| Immunogen                         | Purified chicken IgG (Fc) fragment.   |   |                           |  |                             |  |  |  |
| Specificity                       | Goat anti Chicken IgG (For<br>epitopes within the Fc region<br>classes as assessed by im<br>does not react with the ligh  | on and sho<br>munoelect   | ws no cros<br>rophoresis. | s-reactivity with other ch<br>This Goat anti Chicken | icken immunoglobulin        |  |  |  |

|                                  | Goat anti chicken IgG (Fc) has been used successfully for the evaluation of circulating levels of IgG in chickens using ELISA in a number of experimental and field situations.   |  |  |  |
|----------------------------------|---|--|--|--|
| References                       | <ol> <li>Norup, L.R. <i>et al.</i> (2009) Influence of chicken serum mannose-binding lectin levels on the immune response towards <i>Escherichia coli</i>. <u>Poult Sci. 88:543-53</u>.</li> <li>Duckworth, J.A. <i>et al.</i> (2008) Development of a contraceptive vaccine for the marsupial brushtail possum (<i>Trichosurus vulpecula</i>): lack of effects in mice and chickens immunised with recombinant possum ZP3 protein and a possum ZP3 antifertility epitope <u>Wildlife Research 35, 563–72</u>.</li> <li>Pleidrup, J. <i>et al.</i> (2014) <i>Ascaridia galli</i> infection influences the development of both humoral and cell-mediated immunity after Newcastle Disease vaccination in chickens. <u>Vaccine. 32 (3): 383-92</u>.</li> <li>Peralta, B. <i>et al.</i> (2009) Evidence of widespread infection of avian hepatitis E virus (avian HEV) in chickens from Spain. <u>Vet Microbiol. 137: 31-6</u></li> <li>Cho Y <i>et al.</i> (2015) Proteomic analysis of outer membrane proteins in <i>Salmonella enterica Enteritidis</i>. J Microbiol Biotechnol. 25 (2): 288-95.</li> <li>Radomska KA <i>et al.</i> (2016) Chicken Immune Response after <i>In Ovo</i> Immunization with Chimeric TLR5 Activating Flagellin of <i>Campylobacter jejuni</i>. <u>PLoS One. 11 (10): e0164837</u>.</li> <li>Vaezirad, M.M. <i>et al.</i> (2018) Chicken immune response following in ovo delivery of bacterial flagellin. <u>Vaccine. Mar 09 [Epub ahead of print].</u></li> </ol> |  |  |  |
| Storage                          | Store at +4°C. DO NOT FREEZE.<br>This product should be stored undiluted. Should this product contain a precipitate we recommend<br>microcentrifugation before use.   |  |  |  |
| Shelf Life                       | 12 months from date of despatch.  |  |  |  |
| Acknowledgements                 | Proclin™ 300 is a trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow.  |  |  |  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #20391 available at: 20391: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/20391.pdf</u>  |  |  |  |
| Regulatory                       | For research purposes only  |  |  |  |

### **Related Products**

### **Recommended Useful Reagents**

AbGUARD® HRP STABILIZER PLUS (BUF052A) AbGUARD® HRP STABILIZER PLUS (BUF052B) AbGUARD® HRP STABILIZER PLUS (BUF052C) TMB CORE (BUF056A) TMB CORE+ (BUF062A) TMB SIGNAL+ (BUF054A)

| North & South | Tel: +1 800 265 7376        | Worldwide | Tel: +44 (0)1865 852 700      | Europe  | Tel: +49 (0) 89 8090 95 21           |
|---------------|-----------------------------|-----------|-------------------------------|---------|--------------------------------------|
| America       | Fax: +1 919 878 3751        |           | Fax: +44 (0)1865 852 739      |         | Fax: +49 (0) 89 8090 95 50           |
|               | Email: antibody_sales_us@bi | o-rad.com | Email: antibody_sales_uk@bio- | rad.com | Email: antibody_sales_de@bio-rad.com |

'M314312:180412'

#### Printed on 02 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint