

## Datasheet: AAI34

<b>Description:</b>	GOAT ANTI HORSE IgA
<b>Specificity:</b>	IgA
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/100 - 1/1000
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Horse
<b>Product Form</b>	Purified IgG - liquid
<b>Antiserum Preparation</b>	Antisera to equine IgA were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography using antigen coupled to agarose beads.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified equine IgA
<b>Specificity</b>	<b>Goat anti Horse IgA antibody</b> recognizes equine IgA and shows no cross-reactivity with other equine immunoglobulin classes in immunoelectrophoresis. Goat anti Horse IgA antibody may cross-react with IgA from other species.

**References**

1. Lewis, M.J. *et al.* (2010) IgA in the horse: cloning of equine polymeric Ig receptor and J chain and characterization of recombinant forms of equine IgA. [Mucosal Immunol. 3: 610-21.](#)
2. Keggan, A. *et al.* (2013) Production of seven monoclonal equine immunoglobulins isotyped by multiplex analysis. [Vet Immunol Immunopathol. 153 \(3-4\): 187-93.](#)
3. van der Kolk, J.H. *et al.* (2012) Gluten-dependent antibodies in horses with inflammatory small bowel disease (ISBD). [Vet Q. 32: 3-11.](#)

---

**Storage**

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Shelf Life**

18 months from date of despatch.

---

**Health And Safety Information**

Material Safety Datasheet documentation available at:  
Material Safety Datasheet Documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

**North & South America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M314110:180412'

**Printed on 13 Apr 2018**