

## Datasheet: AAI38

<b>Description:</b>	GOAT ANTI HORSE IgG (T)
<b>Specificity:</b>	IgG (T)
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA	■			1/100 - 1/1000
Immunoprecipitation			■	
Western Blotting			■	
Immunodiffusion	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Horse
<b>Product Form</b>	Purified IgG - liquid
<b>Antiserum Preparation</b>	Antisera to equine IgG (T) were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography using antigen coupled to agarose beads.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified equine IgG (T).
<b>Specificity</b>	<b>Goat anti Horse IgG (T) antibody</b> recognizes equine IgG (T). No cross-reactivity with other equine immunoglobulin classes is seen in immuno-electrophoresis.

Goat anti Horse IgG (T) antibody may cross react with IgG from other species.

---

## References

1. Hooper-McGrevy, K.E. *et al.* (2003) Immunoglobulin G subisotype responses of pneumonic and healthy, exposed foals and adult horses to *Rhodococcus equi* virulence-associated proteins. [Clin Diagn Lab Immunol. 10 \(3\): 345-51.](#)
2. Jacks, S. *et al.* (2007) Experimental infection of neonatal foals with *Rhodococcus equi* triggers adult-like gamma interferon induction. [Clin Vaccine Immunol. 14: 669-77.](#)
3. Lewis, M.J. *et al.* (2007) The different effector function capabilities of the seven equine IgG subclasses have implications for vaccine strategies. [Mol Immunol. 45: 818-27.](#)
4. Ryan, C. & Giguère, S. (2010) Equine neonates have attenuated humoral and cell-mediated immune responses to a killed adjuvanted vaccine compared to adult horses. [Clin Vaccine Immunol. 17 \(12\): 1896-902.](#)
5. Cauchard S *et al.* (2014) Assessment of the safety and immunogenicity of *Rhodococcus equi*-secreted proteins combined with either a liquid nanoparticle (IMS 3012) or a polymeric (PET GEL A) water-based adjuvant in adult horses and foals--identification of promising new candidate antigens. [Vet Immunol Immunopathol. 157 \(3-4\): 164-74.](#)
6. Meulenbroeks C *et al.* (2015) Allergen-Specific Cytokine Polarization Protects Shetland Ponies against *Culicoides obsoletus*-Induced Insect Bite Hypersensitivity. [PLoS One. 10 \(4\): e0122090.](#)
7. Cauchard, S. *et al.* (2014) Assessment of the safety and immunogenicity of *Rhodococcus equi*-secreted proteins combined with either a liquid nanoparticle (IMS 3012) or a polymeric (PET GEL A) water-based adjuvant in adult horses and foals--identification of promising new candidate antigens. [Vet Immunol Immunopathol. 157 \(3-4\): 164-74.](#)
8. Burk, S.V. *et al.* (2016) Equine antibody response to larval *Parascaris equorum* excretory-secretory products. [Vet Parasitol. 226: 83-7.](#)
9. Lightbody, K.L. *et al.* (2016) Validation of a novel saliva-based ELISA test for diagnosing tapeworm burden in horses. [Vet Clin Pathol. 45 \(2\): 335-46.](#)

---

## Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

---

## Shelf Life

18 months from date of despatch.

---

## Health And Safety Information

Material Safety Datasheet documentation #10040 available at:  
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

## Regulatory

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M314433:180412'

Printed on 01 May 2018

---