

Datasheet: AAI41B

Description:	GOAT ANTI PIG IgG (Fc):Biotin
Specificity:	IgG (Fc)
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1:10000 - 1:100000
Western Blotting	▪			1:10000 - 1:100000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig
Product Form	Purified IgG fraction conjugated to Biotin - liquid
Antiserum Preparation	Antisera to porcine IgG were raised by repeated immunisation of goat with highly purified antigen. Purified IgG was prepared by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	0.2% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified porcine IgG.

Specificity

Goat anti Pig IgG (Fc) antibody recognizes the Fc region of the porcine IgG heavy chains and shows no cross - reactivity with other porcine immunoglobulin classes as evaluated by immunoelectrophoresis. Goat anti Pig IgG (Fc) has not been species cross adsorbed and may react with the Fc region of IgG from other species.

Goat anti Pig IgG (Fc) antibody has been used extensively as a detection reagent for porcine IgG in ELISA, for example monitoring of the IgG response in influenza infected pigs ([Crisci et al. 2013](#)).

References

1. Assana, E. *et al.* (2010) Antibody responses to the host-protective *Taenia solium* oncosphere protein TSOL18 in pigs are directed against conformational epitopes. [Parasite Immunol. 32: 399-405.](#)
2. Kick, A.R. *et al.* (2011) Evaluation of peripheral lymphocytes after weaning and vaccination for *Mycoplasma hyopneumoniae*. [Res Vet Sci. 91 \(3\): e68-72.](#)
3. Pyo, H. *et al.* (2010) Serodiagnosis of porcine reproductive and respiratory syndrome virus infection with the use of glycoprotein 5 antigens. [Can J Vet Res. 74: 223-7.](#)
4. Busquets, N. *et al.* (2010) Experimental infection with H1N1 European swine influenza virus protects pigs from an infection with the 2009 pandemic H1N1 human influenza virus. [Vet Res. 41: 74.](#)
5. Kang, M.L. *et al.* (2008) Chitosan microspheres containing *Bordetella bronchiseptica* antigens as novel vaccine against atrophic rhinitis in pigs. [J Microbiol Biotechnol. 18: 1179-85.](#)
6. Kim, T.*et al.* (2009) *Bordetella bronchiseptica* aroA mutant as a live vaccine vehicle for heterologous porcine circovirus type 2 major capsid protein expression. [Vet Microbiol. 138: 318-24.](#)
7. Picherot, M. *et al.* (2007) Swine infection with *Trichinella spiralis*: Comparative analysis of the mucosal intestinal and systemic immune responses. [Vet Parasitol. 143: 122-30.](#)
8. Tsai, Y.C. *et al.* (2010) Porcine circovirus type 2 (PCV2) induces cell proliferation, fusion, and chemokine expression in swine monocytic cells *in vitro*. [Vet Res. 41: 60.](#)
9. Scharek, L. *et al.* (2005) Influence of a probiotic *Enterococcus faecium* strain on development of the immune system of sows and piglets. [Vet Immunol Immunopathol. 105: 151-61.](#)
10. Scharek, L. *et al.* (2007) Impact of the probiotic bacteria *Enterococcus faecium* NCIMB 10415 (SF68) and *Bacillus cereus* var. *toyoi* NCIMB 40112 on the development of serum IgG and faecal IgA of sows and their piglets. [Arch Anim Nutr. 61: 223-34.](#)
11. Sheoran, A. *et al.* (2012) Infection With *Cryptosporidium hominis* Provides Incomplete Protection of the Host Against *Cryptosporidium parvum*. [J Infect Dis 205: 1019-23.](#)
12. Pina-Pedrero, S. *et al.* (2012) Genomic and antigenic characterization of monomeric autotransporters of *Haemophilus parasuis*: an ongoing process of reductive evolution. [Microbiology. 158: 436-47.](#)
13. Fleury, A. *et al.* (2015) *Taenia solium*: Development of an Experimental Model of Porcine Neurocysticercosis. [PLoS Negl Trop Dis. 9 \(8\): e0003980.](#)
14. Zhang Y *et al.* (2016) Generation of *E. coli*-derived virus-like particles of porcine circovirus type 2 and their use in an indirect IgG enzyme-linked immunosorbent assay. [Arch Virol. Mar 14. \[Epub ahead of print\]](#)
15. Lee, J-A. *et al.* (2016) *Mycoplasma hyorhinis* is a potential pathogen of porcine respiratory disease complex that aggravates pneumonia caused by porcine reproductive and respiratory syndrome virus [Vet Immunol immunopathol. Jun 16 \[Epub ahead of print\]](#)
16. Lorenzen E *et al.* (2017) Intrauterine inoculation of minipigs with *Chlamydia trachomatis* during diestrus establishes a longer lasting infection compared to vaginal inoculation during estrus. [Microbes Infect. Feb 9. pii: S1286-4579\(17\)30025-4. \[Epub ahead of print\]](#)

Storage

Store at +4°C. DO NOT FREEZE.
This product should be stored undiluted.

Shelf Life

12 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com
'M314225:180412'

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 11 Jul 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)