

Datasheet: AHP031F

| Description: | SHEEP ANTI HUMAN C3c:FITC |
|---------------|---------------------------|
| Specificity: | C3c |
| Other names: | COMPLEMENT COMPONENT 3c |
| Format: | FITC |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 1 ml |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | • | |
| Immunohistology - Frozen | • | | | 1/50 - 1/100 |
| Immunohistology - Paraffin | | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| Target Species | Human |
|----------------|-------|
|----------------|-------|

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
|-----------|-------------|---------------------|-------------------|
| | FITC | 490 | 525 |

Antiserum Preparation Antisera to human C3c were raised by repeated immunisation of sheep with highly purified antigen. Purified IgG was prepared by ion exchange chromatography.

| Buffer Solution | Phosphate buffered saline |
|-----------------------------------|-------------------------------|
| Preservative Stabilisers | 0.09% Sodium Azide |
| Approx. Protein Concentrations | IgG concentration 10 mg/ml |
| Immunogen | Human C3 purified from serum. |
| | |

External Database

Links UniProt:

P01024 Related reagents

Entrez Gene:

718 C3 Related reagents

| Synonyms | CPAMD1 |
|--------------------------------------|---|
| Specificity | Sheep anti Human C3c antibody recognizes the C3c component of human complement, formed as a result of the inactivation of C3b. Sheep anti Human C3c antibody may be used for the detection of C3 deposits in tissues following complement activation. |
| Histology Positive Control Tissue | Human skin |
| References | 1. Kennedy, M.W. & Kuo, Y.M. (1988) The surfaces of the parasitic nematodes <i>Trichinella spiralis</i> and <i>Toxocara canis</i> differ in the binding of post-C3 components of human complement by the alternative pathway. Parasite Immunol. 10:459-63. 2. Oyeyinka, G.O. et al. (2003) The effects of ageing on the immune response to <i>Schistosoma haematobium</i> and hookworm by measuring circulating immune complexes, C3, IgG, IgA and IgM levels in residents of Omi dam area of Kogi State, Nigeria. Afr J Med Med Sci. 32: 263-7. 3. Rojana-Udomsart, A. et al. (2013) Complement-mediated muscle cell lysis: A possible mechanism of myonecrosis in anti-SRP associated necrotizing myopathy (ASANM). J Neuroimmunol. 264: 65-70. 4. Akinlade, K.S. et al. (2004) Circulating immune complexes, immunoglobulin classes (IgG, IgA and IgM) and complement components (C3c, C4 and Factor B) in diabetic Nigerians. West Afr J Med. 23 (3): 253-5. |
| Storage | Store at +4°C or at -20°C if preferred. |
| | This product should be stored undiluted. |
| | Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. |
| | Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |
| Shelf Life | 18 months from date of despatch. |
| Health And Safety | Material Safety Datasheet documentation #10040 available at: |
| Information | 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf |

America

North & South Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301222:170109'

Printed on 10 May 2018