

Datasheet: AHP2318

Description:	RABBIT ANTI CD54 / ICAM-1
Specificity:	CD54 / ICAM-1
Other names:	ICAM
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			0.5 - 4.0 ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	30% Glycerol 0.5% Bovine Serum Albumin 0.01% Thiomersal
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Synthetic peptide surrounding amino acid 359 of human CD54 / ICAM-1
External Database Links	UniProt: P05362 Related reagents Entrez Gene: 3383 ICAM1 Related reagents

Specificity Rabbit anti CD54 / ICAM-1 antibody recognizes the CD54 antigen also known as intercellular adhesion molecule 1 (ICAM-1). CD54 is a 532 amino acid single pass type I membrane glycoprotein containing five [Ig-like C2-type domains](#). CD54 is widely expressed and acts as a ligand for the leukocyte integrin LFA-1 (CD11a/CD18) and Mac-1 integrins, it is the major receptor for human rhinovirus ([Staunton et al. 1989](#)).

CD54 is highly expressed on activated endothelial cells ([Favoloro 1993](#)) and plays a role in transendothelial cell migration to sites of acute inflammation, functioning primarily as an adhesion molecule between endothelial cells and leucocytes, following injury or stress ([Manes and Pober 2011](#)). Levels of soluble CD54 in serum ([Matalliotakis et al. 2001](#)) or urine ([Kim et al. 2006](#)) can be potentially used as an indication of infection, inflammation and certain tumors, and as a target for immunosuppression in transplantation.

Western Blotting Rabbit anti CD54 / ICAM-1 antibody recognizes a band of approximately 89-92 kDa in rat kidney tissue lysate, 3T3 cell lysate and Jurkat cell lysate

Storage Store at -20°C
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10096 available at:
10096: <https://www.bio-rad-antibodies.com/uploads/MSDS/10096.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
Sheep Anti Rabbit IgG (2AB02...) [Biotin](#)
Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®549](#), [DyLight®649](#),
[DyLight®680](#), [DyLight®800](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M293176:160810'

Printed on 01 May 2018