Datasheet: AHP2581

Description:	RABBIT ANTI C-JUN (pTyr170)
Specificity:	C-JUN (pTyr170)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/500 - 1/2000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Product Form

Purified IgG - liquid

Antiserum Preparation

Antiserum to c-Jun (pTyr170) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.02% Sodium Azide
50% Glycerol

Immunogen

Phospho specific-peptide corresponding to residues surrounding tyrosine 170 of human c-Jun

External Database Links

UniProt:
P05412 Related reagents

Entrez Gene:
3725 JUN Related reagents

Specificity

Rabbit anti c-Jun (pTyr170) antibody recognizes proto-oncogene c-Jun, also known as V-jun avian sarcoma virus 17 oncogene homolog and p39, when phosphorylated at tyrosine 170. c-Jun is a member of the Jun family of transcription factors and is a subunit of the AP-1 transcription factor. c-Jun dimerizes with c-Fos, JDP and ATF family members to form the AP-1 transcription factor. AP-1 is activated by a variety of stimuli including cytokines, stress and growth factors and plays a

	critical role in cellular processes ranging from proliferation to apoptosis. c-Jun is phosphorylated at tyrosine 170 by the c-Abl tyrosine kinase (Gao *et al.* 2006) preventing ubiquitin mediated degradation of c-Jun.
Western Blotting	Due to the presence of phosphoproteins in milk, the use of milk based blocking reagents is not recommended. 1% BSA in PBS or TBS Tween should be used instead.
Further Reading	1. Gao, B. *et al.* (2006) The tyrosine kinase c-Abl protects c-Jun from ubiquitination-mediated degradation in T cells. J Biol Chem. 281 (40): 29711-8.
Storage	Store at -20^{o}C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: 10049: https://www.bio-rad-antibodies.com/uploads/MSDS/10049.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)	FITC
Sheep Anti Rabbit IgG (STAR35...)	RPE
Goat Anti Rabbit IgG (H/L) (STAR124...)	HRP
Goat Anti Rabbit IgG (Fc) (STAR121...)	Biotin, FITC, HRP
Sheep Anti Rabbit IgG (2AB02...)	Biotin
Sheep Anti Rabbit IgG (STAR36...)	DyLight®488, DyLight®549, DyLight®649, DyLight®680, DyLight®800

Recommended Useful Reagents

TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South America Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M303352:170220'

Printed on 01 May 2018