

Datasheet: AHP2586

Description:	RABBIT ANTI EGF R (pSer1046)
Specificity:	EGF R (pSer1046)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/500 - 1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG - liquid

Antiserum Preparation Antiserum to EGF R (pSer1046) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution Phosphate buffered saline

Preservative 0.02% Sodium Azide
Stabilisers 50% Glycerol

Immunogen Phospho specific-peptide corresponding to residues surrounding serine1046 of human EGF R

External Database Links

UniProt:

[P00533](#) [Related reagents](#)

Entrez Gene:

[1956](#) EGFR [Related reagents](#)

Synonyms ERBB1

Specificity **Rabbit anti EGF R (pSer1046) antibody** recognizes the epidermal growth factor receptor (EGF R), also known as proto-oncogene c-ErbB-1 or receptor tyrosine-protein kinase erbB-1, when phosphorylated at serine 1046. EGF R is a transmembrane glycoprotein and a member of the ErbB

receptor tyrosine kinase family. EGFR is a cell surface receptor for members of the epidermal growth factor family.

EGF R plays a role in numerous cellular processes such as cell proliferation and apoptosis through the activation of downstream signaling pathways ([Scaltriti and Baselga 2006](#)).

EGF R is autophosphorylated at serine 1046 and serine 1047 upon ligand binding and receptor dimerization. Phosphorylation of serine 1046 and serine 1047 plays an important role in EGF R desensitization ([Sun et al. 2010](#)).

Western Blotting	Due to the presence of phosphoproteins in milk, the use of milk based blocking reagents is not recommended. 1% BSA in PBS or TBS Tween should be used instead.
Further Reading	1. Scaltriti, M. & Baselga, J. (2006) The epidermal growth factor receptor pathway: a model for targeted therapy. Clin Cancer Res. 12 (18): 5268-72. 2. Sun, Q. <i>et al.</i> (2010) Additional serine/threonine phosphorylation reduces binding affinity but preserves interface topography of substrate proteins to the c-Cbl TKB domain. PLoS One. 5 (9): e12819.
Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: 10049: https://www.bio-rad-antibodies.com/uploads/MSDS/10049.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
Sheep Anti Rabbit IgG (2AB02...) [Biotin](#)
Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®549](#), [DyLight®649](#),
[DyLight®680](#), [DyLight®800](#)

Recommended Useful Reagents

[TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M303353:170220'

