

Datasheet: AHP2599

Description:	RABBIT ANTI ELK-1 (pSer383)
Specificity:	ELK-1 (pSer383)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin	■			1/50 - 1/100
Western Blotting	■			1/500 - 1/2000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Based on sequence similarity, is expected to react with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Antiserum Preparation	Antiserum to ELK-1 (pSer383) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative	0.02% Sodium Azide
Stabilisers	50% Glycerol
Immunogen	Phospho specific-peptide corresponding to residues surrounding serine 383 of human ELK-1
External Database Links	UniProt: P19419 Related reagents Entrez Gene: 2002 ELK1 Related reagents
Specificity	Rabbit anti ELK-1 (pSer383) antibody recognizes ELK-1, also known as ETS domain-containing

protein Elk-1, when phosphorylated at serine 383.

ELK-1 is a member of the ternary complex factor (TCF) family of transcription factors, and plays a role in mediating gene expression in response to growth factors.

ELK-1 is phosphorylated at serine 383 and 389 in its C-terminal region by mitogen activated protein (MAP) kinase. Phosphorylation of serine 383 and 389 is important for activation of ELK1 ([Li et al. 2003](#)).

Western Blotting	Due to the presence of phosphoproteins in milk, the use of milk based blocking reagents is not recommended. 1% BSA in PBS or TBS Tween should be used instead.
Further Reading	1. Li, Q.J. <i>et al.</i> (2003) MAP kinase phosphorylation-dependent activation of Elk-1 leads to activation of the co-activator p300. EMBO J. 22 (2): 281-91.
Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: 10049: https://www.bio-rad-antibodies.com/uploads/MSDS/10049.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)	FITC
Sheep Anti Rabbit IgG (STAR35...)	RPE
Goat Anti Rabbit IgG (H/L) (STAR124...)	HRP
Goat Anti Rabbit IgG (Fc) (STAR121...)	Biotin , FITC , HRP
Sheep Anti Rabbit IgG (2AB02...)	Biotin
Sheep Anti Rabbit IgG (STAR36...)	DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800

Recommended Useful Reagents

[TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M303465:170222'

Printed on 01 May 2018