

Datasheet: AHP2615

Description:	RABBIT ANTI FOCAL ADHESION KINASE (pTyr576/pTyr577)
Specificity:	FOCAL ADHESION KINASE (pTyr576/pTyr577)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/500 - 1/2000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Antiserum Preparation	Antiserum to focal adhesion kinase (pTyr576/pTyr577) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.02% Sodium Azide 50% Glycerol
Immunogen	Phospho specific-peptide corresponding to residues surrounding tyrosine 576/ tyrosine 577 of human focal adhesion kinase
External Database Links	UniProt: Q05397 Related reagents Entrez Gene: 5747 PTK2 Related reagents
Synonyms	FAK, FAK1
Specificity	Rabbit anti focal adhesion kinase (pTyr576/pTyr577) antibody recognizes focal adhesion

kinase, also known as FAK, when phosphorylated at tyrosine 576/ tyrosine 577. Focal adhesion kinase is a highly conserved tyrosine kinase with roles in several biological processes including adhesion, migration and cell survival. FAK consists of four domains, an N-terminal FERM domain; a catalytic tyrosine kinase domain; a C-terminal focal-adhesion targeting (FAT) domain and an unstructured proline rich region between the catalytic and FAT domains. Phosphorylation of FAK at tyrosine 576/577 by Src leads to an increase in the kinase activity of FAK.

Western Blotting	Due to the presence of phosphoproteins in milk, the use of milk based blocking reagents is not recommended. 1% BSA in PBS or TBS Tween should be used instead.
Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: 10049: https://www.bio-rad-antibodies.com/uploads/MSDS/10049.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
 Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
 Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
 Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
 Sheep Anti Rabbit IgG (2AB02...) [Biotin](#)
 Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®549](#), [DyLight®649](#),
[DyLight®680](#), [DyLight®800](#)

Recommended Useful Reagents

[TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M303502:170222'

Printed on 01 May 2018