

Datasheet: AHP669

Description:	RABBIT ANTI GAP43
Specificity:	GAP43
Format:	Serum
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			1/500 - 1/1000
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Cat
Species Cross Reactivity	Reacts with: Human, Rat, Monkey, Mouse N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Serum - liquid

Antiserum Preparation Antisera to GAP43 were raised by repeated immunisations of rabbits with highly purified antigen.

Preservative Stabilisers	0.02% Sodium Azide (NaN ₃)
---------------------------------	--

Immunogen	Purified GAP43 from cat brain.
------------------	--------------------------------

External Database Links	UniProt: Q6S9D9 Related reagents
	Entrez Gene: 493873 GAP43 Related reagents

Specificity **Rabbit anti GAP43 antibody** recognises the protein designated GAP43, a neuronal protein that is expressed at elevated levels during development and also during regenerative axon growth.

GAP43 (also known as neuromodulin, pp46, B-50, Protein F1 and calmodulin-binding protein P-57) is a protein of approximately 25 kDa that migrates on SDS-PAGE gels with an apparent size of 40-50 kDa. The aberrant migration is due to the unusually low abundance of hydrophobic amino acid residues. GAP43 is found tightly associated with neuronal membranes due to palmitoylation. The solubilised protein binds calmodulin, although this is reduced following protein kinase C phosphorylation of GAP43.

References

1. McIntosh, H. *et al.* (1989) A GAP-43-like protein in cat visual cortex. [Vis Neurosci. 2 \(6\): 583-91.](#)
2. Wang, H. *et al.* (2009) Neuroprotective effect of recombinant human erythropoietin on optic nerve injury in rats. [Chin Med J 122: 2008-12.](#)
3. Zhang, H. *et al.* (2010) Alteration of parasympathetic/sympathetic ratio in the infarcted myocardium after Schwann cell transplantation modified electrophysiological function of heart: a novel antiarrhythmic therapy. [Circulation. 122 \(11 Suppl\): S193-200.](#)
4. Parikh, P. *et al.* (2011) Regeneration of axons in injured spinal cord by activation of bone morphogenetic protein/Smad1 signaling pathway in adult neurons. [Proc Natl Acad Sci U S A. 108: E99-107.](#)
5. Turner, J.H. *et al.* (2010) Reversible loss of neuronal marker protein expression in a transgenic mouse model for sinusitis-associated olfactory dysfunction. [Am J Rhinol Allergy. 24: 192-6.](#)
6. Su, Y. *et al.* (2008) Axonal regeneration after optic nerve crush in Nogo-A/B/C knockout mice. [Mol Vis. 14: 268-73.](#)
7. Pao, P.C. *et al.* (2011) A novel RING finger protein, Znf179, modulates cell cycle exit and neuronal differentiation of P19 embryonal carcinoma cells. [Cell Death Differ. 18 \(11\): 1791-804.](#)
8. Cheng, H.T. *et al.* (2013) Increased Axonal Regeneration and Swellings in Intraepidermal Nerve Fibers Characterize Painful Phenotypes of Diabetic Neuropathy. [J Pain. pii: S1526-5900\(13\)00910-3.](#)
9. Wu, X. *et al.* (2012) Desipramine pretreatment improves sympathetic remodeling and ventricular fibrillation threshold after myocardial ischemia. [J Biomed Biotechnol. 2012: 732909.](#)
10. Oberbauer, E. *et al.* (2013) Chroman-like cyclic prenylflavonoids promote neuronal differentiation and neurite outgrowth and are neuroprotective. [J Nutr Biochem. 24 \(11\): 1953-62.](#)
11. Su, Y. *et al.* (2017) Transfection with AAV-NgR1siRNA on axonal regeneration of optic nerve after injury [Int J Clin Exp Pathol. 10 \(2\) :2327-32.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10081 available at:
10081: <https://www.bio-rad-antibodies.com/uploads/MSDS/10081.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
Sheep Anti Rabbit IgG (2AB02...) [Biotin](#)
Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®549](#), [DyLight®649](#),
[DyLight®680](#), [DyLight®800](#)

Recommended Useful Reagents

[TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M317057:180618'

Printed on 21 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)