

Datasheet: AHP999

Description:	RABBIT ANTI MOUSE ST2 (N-TERMINAL)
Specificity:	ST2 (N-TERMINAL)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting	■			1 - 2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Antiserum Preparation	Antisera to mouse ST2 (NT) were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.02% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	A 16 amino acid synthetic peptide located near mouse ST2 amino-terminus. This peptide is common to the three known isoforms of ST2.

External Database Links

UniProt:

[P14719](#)

[Related reagents](#)

Entrez Gene:[17082](#) [Il1rl1](#) [Related reagents](#)

Synonyms	Ly84, St2, Ste2
-----------------	-----------------

Specificity	<p>Rabbit anti Mouse ST2 (N-Terminal) antibody recognizes an epitope within the N-Terminal region (NT) of mouse ST2, a member of the Toll-interleukin 1 receptor (TIR) superfamily, expressed by mast cells and TH₂ helper cells, but not by TH₁ effector cells.</p> <p>The majority of TIR superfamily members, which includes the Interleukin-1 receptor (IL-1R1), Toll-like receptors (TLRs) and the cytosolic adaptor proteins MyD88, TIRAP, Trif and TRAM, initiate innate immunity through the ultimate activation of the transcription factor NF-kappaB and are inducers of the inflammatory response. In contrast ST2, which itself is unable to activate NF-kappaB, has been identified as a negative regulator of IL-1R1 and TLR4 signaling by sequestering MyD88 and TIRAP in line with its role as a crucial receptor in TH₂-cell mediated responses (Brint <i>et al.</i> 2004 and Coyle <i>et al.</i> 1999).</p>
--------------------	---

Western Blotting	AHP999 detects a band of approximately 45kDa in mouse kidney cell lysates. (Predicted Mwt. 62 kDa).
-------------------------	---

Further Reading	<ol style="list-style-type: none">1. Brint, E.K. <i>et al.</i> (2004) ST2 is an inhibitor of interleukin 1 receptor and Toll-like receptor 4 signaling and maintains endotoxin tolerance. Nat Immunol. 5 (4): 373-9.2. Coyle, A.J. <i>et al.</i> (1999) Crucial role of the interleukin 1 receptor family member T1/ST2 in T helper cell type 2-mediated lung mucosal immune responses. J Exp Med. 190 (7): 895-902.
------------------------	---

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)	FITC
Sheep Anti Rabbit IgG (STAR35...)	RPE
Goat Anti Rabbit IgG (H/L) (STAR124...)	HRP
Goat Anti Rabbit IgG (Fc) (STAR121...)	Biotin , FITC , HRP
Sheep Anti Rabbit IgG (2AB02...)	Biotin
Sheep Anti Rabbit IgG (STAR36...)	DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800

Recommended Useful Reagents

TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M310191:170914'

Printed on 02 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)