

Datasheet: C07SA

Description:	GOAT SERUM
Name:	GOAT SERUM
Format:	Serum
Product Type:	Serum & Cells
Quantity:	100 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	■			
Immunohistology - Paraffin	■			
Immunoassay	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Form Serum - liquid

Antiserum Preparation Whole serum is supplied 0.2µm filtered.

Preservative Stabilisers 0.1% Sodium Azide

References

1. Bürli, T. *et al.* (2010) Single particle tracking of alpha7 nicotinic AChR in hippocampal neurons reveals regulated confinement at glutamatergic and GABAergic perisynaptic sites. [PLoS One. 9;5\(7\):e11507.](https://doi.org/10.1371/journal.plosone.011507)
2. Säve, S. *et al.* (2011) Activation of Adenosine A2A Receptors Inhibits Neutrophil Transuroepithelial Migration. [Infect Immun. 79: 3431-7.](https://doi.org/10.1186/1745-6216-79-3431-7)
3. Luhmann, U.F. *et al.* (2014) The severity of retinal pathology in homozygous Crb1rd8/rd8 mice is dependent on additional genetic factors. [Hum Mol Genet. pii: ddu424. \[](https://doi.org/10.1093/hmg/ddu424)
4. Hippert C *et al.* (2015) Müller glia activation in response to inherited retinal degeneration is highly varied and disease-specific. [PLoS One. 10 \(3\): e0120415.](https://doi.org/10.1371/journal.plosone.0120415)
5. McArthur, S. *et al.* (2015) Sex-specific disruption of murine midbrain astrocytic and dopaminergic developmental trajectories following antenatal GC treatment. [Brain Struct Funct. May 6. \[Epub ahead of print\]](https://doi.org/10.1186/s13041-015-0161-1)
6. Herrmann, P. *et al.* (2015) Cd59a deficiency in mice leads to preferential innate immune activation in the retinal pigment epithelium-choroid with age. [Neurobiol Aging. 36 \(9\): 2637-48.](https://doi.org/10.1007/s12074-015-9488-8)
7. Díaz-Carballo D *et al.* (2015) Therapeutic potential of antiviral drugs targeting chemorefractory colorectal adenocarcinoma cells overexpressing endogenous retroviral elements. [J Exp Clin Cancer Res. 34: 81.](https://doi.org/10.1007/s12074-015-9488-8)

Storage Store at +4°C or at -20°C if preferred.

Storage in frost free freezers is not recommended.

Avoid repeated freezing and thawing as this may denature the product. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 12 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10496 available at:
10496: <https://www.bio-rad-antibodies.com/uploads/MSDS/10496.pdf>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M318957:180719'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)