

Datasheet: DC003

Description:	MOUSE ANTI HUMAN CD3:FITC/CD8:RPE
Specificity:	CD3/CD8
Format:	FITC/RPE
Product Type:	Dual Color Reagent
Clone:	UCHT1/LT8
lsotype:	Cocktail
Quantity:	50 TESTS

Product Details

Applications	from testing within our the originators. Please	laboratories, peer-le refer to references	eviewed publications or indicated for further inf	ons. This information is derived personal communications from ormation. For general protocol	
	recommendations, plea		ad-antibodies.com/proto No Not Determir		
	Flow Cytometry	Tes		ned Suggested Dilution Neat	
	Where this antibody has not been tested for use in a particular technique this does not necessarily				
	exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is				
	recommended that the user titrates the antibody for use in their own system using appropriate				
	negative/positive contr	ols.	-		
Antibody Isotypes	FITC reagent: IgG1 (MOUSE)				
	RPE Reagent: IgG1 (N	MOUSE)			
Target Species	Human				
Product Form	Dual Colour combination consisting of FITC conjugated and RPE conjugated monoclonal antibodies mixed in optimal ratio - lyophilised.				
Reconstitution	Reconstitute with 0.5 ml distilled water				
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)		
	FITC	490	525		
	RPE 488nm laser	496	578		
	RPE 561nm laser	546	578		
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide				
Stabilisers	1% Bovine Serum Albumin				
	5% Sucrose				
External Database Links	UniProt: P04234 Related	<u>d reagents</u>			

	P07766 Related reagents		
	P09693 Related reagents		
	P01732 Related reagents		
	P10966 Related reagents		
	Entrez Gene:		
	915 CD3D Related reagents		
	916 CD3E Related reagents		
	917 CD3G Related reagents		
	925 CD8A Related reagents		
	926 CD8B Related reagents		
Synonyms	CD8B1, MAL, T3D, T3E, T3G		
Specificity	Mouse anti Human CD3:FITC/CD8:RPE dual color reagent may be used to simultaneously identify CD3 and CD8 positive cells and to identify populations of lymphocytes that co-express these antigens.		
	Clone UCHT1 recognises the human CD3 antigen, which is expressed by all mature T cells, while clone LT8 recognises the human CD8 antigen, a ~32 kDa dimer expressed by a subset of T cells with cytotoxic/suppressor activity, and more weakly by NK cells.		
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.		
References	1. Beverley, P. <i>et al.</i> (1981) Distinctive functional characteristics of human "T" lymphocytes defined by E rosetting or a monoclonal anti-T cell antibody. <u>Eur. J. Immunol. 11: 329-334.</u>		
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.		
	This product should be stored undiluted.		
	DO NOT FREEZE. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.		
Shelf Life	12 months from date of reconstitution.		
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</u>		
Regulatory	For research purposes only		

Related Products

Recommended Negative Controls

MOUSE IgG1:FITC/MOUSE IgG1:RPE NEGATIVE CONTROL (DC012)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B) North & South Tel: +1 800 265 7376 America

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com 'M308859:170816'

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Printed on 01 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com

Fax: +1 919 878 3751