

Datasheet: HCA139

| Description: | HUMAN ANTI CHICKEN HEN EGG LYSOZYME |
|---------------|-------------------------------------|
| Specificity: | HEN EGG LYSOZYME |
| Other names: | HEL |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | AbD11397 |
| Isotype: | HuCAL Fab monovalent |
| Quantity: | 0.1 mg |
| | |

Product Details

| Applications | This product has been repo | This product has been reported to work in the following applications. This information is derived | | | | | |
|-----------------------------|--|---|----|----------------|--------------------|--|--|
| | from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol | | | | | | |
| | | | | | | | |
| | recommendations, please visit www.bio-rad-antibodies.com/protocols. | | | | | | |
| | | Yes | No | Not Determined | Suggested Dilution | | |
| | Flow Cytometry | | | | | | |
| | Immunohistology - Frozen | | | | | | |
| | Immunohistology - Paraffin | | | • | | | |
| | ELISA | | | • | | | |
| | Immunoprecipitation | | | • | | | |
| | Western Blotting | • | | | 1ug/ml - 0.1µg/ml | | |
| | Functional Assays | | | | | | |
| | Where this product has not been tested for use in a particular technique this does not necessarily | | | | | | |
| | exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is | | | | | | |
| | recommended that the user titrates the product for use in their own system using appropriate | | | | | | |
| | negative/positive controls. | | | | | | |
| | | | | | | | |
| Target Species | Chicken | | | | | | |
| Species Cross Reactivity | Does not react with:Human | | | | | | |
| Product Form | A monovalent human recombinant Fab (lambda light chain) selected from the HuCAL® phage display library, expressed in <i>E. coli.</i> The antibody is tagged with a V5-tag (GSGGAPGKPIPNPLLGLDSTDAP) and a his-tag (HHHHHH) at the C-terminus of the antibody heavy chain. This antibody is supplied lyophilised. | | | | | | |
| Reconstitution | Reconstitute with 1.0ml distilled water | | | | | | |
| Preparation | metal chelate affinity chromatography | | | | | | |
| Buffer Solution | Phosphate buffered saline | | | | | | |
| Preservative | 0.09% Sodium Azide (NaN ₃) | | | | | | |

| Stabilisers | 1% Bovine Serum Albumin |
|-----------------------------------|--|
| Approx. Protein Concentrations | Total protein concentration 0.1mg/ml |
| Immunogen | Lysozyme from chicken egg white. |
| External Database Links | UniProt: <u>P00698</u> <u>Related reagents</u> Entrez Gene: <u>396218</u> LYZ <u>Related reagents</u> |
| Specificity | Human anti Chicken Hen Egg Lysozyme, clone AbD11397, specifically recognises Hen Egg Lysozyme (HEL), known also as muramidase or N-acetylmuramide glycanhydrolase, a 14kDa enzymic protein involved in the destruction of bacteria. Lysozyme damages bacterial cell walls by catalyzing hydrolysis of 1,4-beta-linkages between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-acetyl-D-glucosamine residues in chitodextrins. Lysozyme is abundant in a number of secretions, such as tears, saliva, human milk and mucus. It is abundant in a number of secretions. |
| | is also present in cytoplasmic granules of PMN's and high concentrations of lysozyme are present in egg white. C-type lysozymes are closely related to alpha-lactalbumin in sequence and structure making them part of the same family. |
| ELISA | Due to the presence of bovine serum albumin (BSA), this antibody is unsuitable for use as a capture reagent in sandwich ELISA applications. This product is also available without BSA, <u>please</u> enquire. |
| Storage | Prior to reconstitution store at +4°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |
| Shelf Life | 12 months from date of reconstitution. |
| Acknowledgements | Sold under license of U.S. Patents 6753136, 7785859 and 8273688 and corresponding patents. This antibody was developed by Bio-Rad, Zeppelinstr. 4, 82178 Puchheim, Germany. His-tag is a registered trademark of EMD Biosciences. |
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: 10041: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</u> |
| Regulatory | For research purposes only |
| Technical Advice | Recommended protocols and further information about HuCAL recombinant antibody technology can be found in the HuCAL Antibodies Technical Manual |

Related Products

Recommended Secondary Antibodies

Mouse Anti Synthetic Peptide HISTIDINE TAG (MCA5995...) HRP Goat Anti Human IgG F(ab')2 (0500-0099...) HRP Mouse Anti Synthetic Peptide HISTIDINE TAG (MCA1396...) ALEXA FLUOR® 488, ALEXA FLUOR® 647, Alk. Phos., Biotin, FITC

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M311309:171113'

Printed on 04 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint