

Datasheet: MCA1075BT

Description:	MOUSE ANTI HUMAN CD32:Biotin
Specificity:	CD32
Other names:	FcRII
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	AT10
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Dog, Rhesus Monkey, Pig N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	K562 cell line.
External Database Links	UniProt: P12318 Related reagents

[P31994](#) [Related reagents](#)

[P31995](#) [Related reagents](#)

Entrez Gene:

[2212](#) FCGR2A [Related reagents](#)

[2213](#) FCGR2B [Related reagents](#)

[9103](#) FCGR2C [Related reagents](#)

Synonyms	CD32, FCG2, FCGR2A1, IGFR2
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	Mouse anti Human CD32 antibody, clone AT10 recognizes the human CD32 antigen, a ~40 kDa glycoprotein that acts as a low affinity receptor for IgG (also known as Fc gamma RII). The antigen mediates several functions including endocytosis, activation of secretion, cytotoxicity and immunomodulation. CD32 is expressed by B cells, monocytes, granulocytes and platelets. Clone AT10 blocks the binding of IgG to Fc gamma RII (Larsson et al. 1997).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
References	1. Greenman, J. <i>et al.</i> (1991) Characterization of a new monoclonal anti-Fc gamma RII antibody, AT10, and its incorporation into a bispecific F(ab')₂ derivative for recruitment of cytotoxic effectors. Mol Immunol. 28 (11): 1243-54. 2. Van Den Herik-Oudijk, I.E. <i>et al.</i> (1994) Functional analysis of human Fc gamma RII (CD32) isoforms expressed in B lymphocytes. J Immunol. 152 (2): 574-85. 3. Lilliehöök, I. <i>et al.</i> (1998) Expression of adhesion and Fc gamma-receptors on canine blood eosinophils and neutrophils studied by anti-human monoclonal antibodies. Vet Immunol Immunopathol. 61 (2-4): 181-93. 4. Larsson M <i>et al.</i> (1997) Human dendritic cells handling of binding, uptake and degradation of free and IgG-immune complexed dinitrophenylated human serum albumin <i>in vitro</i>. Immunology. 90 (1): 138-46. 5. Dutertre, C.A. <i>et al.</i> (2008) A novel subset of NK cells expressing high levels of inhibitory Fc gammaRIIB modulating antibody-dependent function. J Leukoc Biol. 84 (6): 1511-20. 6. Devriendt, B. <i>et al.</i> (2010) Targeting of <i>Escherichia coli</i> F4 fimbriae to Fc gamma receptors enhances the maturation of porcine dendritic cells. Vet Immunol Immunopathol. 135: 188-98. 7. Sims, G.P. <i>et al.</i> (2005) Identification and characterization of circulating human transitional B cells. Blood. 105: 4390-8. 8. Benitez-Ribas, D. <i>et al.</i> (2006) Plasmacytoid dendritic cells of melanoma patients present exogenous proteins to CD4+ T cells after Fc gamma RII-mediated uptake. J Exp Med. 203: 1629-35. 9. Zhao, X.W. <i>et al.</i> (2011) CD47-signal regulatory protein-α (SIRPα) interactions form a barrier for antibody-mediated tumor cell destruction. Proc Natl Acad Sci U S A. 108 (45): 18342-7. 10. Araújo, M.S.<i>et al.</i> (2011) Immunological changes in canine peripheral blood leukocytes triggered by immunization with first or second generation vaccines against canine visceral leishmaniasis. Vet Immunol Immunopathol. 141: 64-75. 11. Benítez-Ribas, D. <i>et al.</i> (2006) Plasmacytoid dendritic cells of melanoma patients present exogenous proteins to CD4+ T cells after Fc gamma RII-mediated uptake. J Exp Med. 203: 1629-35. 12. Santer, D.M. <i>et al.</i> (2010) C1q deficiency leads to the defective suppression of IFN-alpha in response to nucleoprotein containing immune complexes. J Immunol. 185: 4738-49.

13. Shannon, O. *et al.* (2010) Platelet activation and biofilm formation by *Aerococcus urinae*, an endocarditis-causing pathogen. [Infect Immun. 78: 4268-75.](#)
14. Ito, T. *et al.* (1999) A CD1a+/CD11c+ subset of human blood dendritic cells is a direct precursor of Langerhans cells. [J Immunol. 163: 1409-19.](#)
15. Moreira, M.L. *et al.* (2016) Vaccination against canine leishmaniosis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. [Vet Parasitol. 220: 33-45.](#)
16. Gazendam, R.P. *et al.* (2016) Impaired killing of *Candida albicans* by granulocytes mobilized for transfusion purposes: a role for granule components. [Haematologica. 101 \(5\): 587-96.](#)
17. Liu M *et al.* (2011) Vitellogenin mediates phagocytosis through interaction with FcγR. [Mol Immunol. 49 \(1-2\): 211-8.](#)
18. Petersson, F. *et al.* (2018) Platelet activation and aggregation by the opportunistic pathogen *Cutibacterium (Propionibacterium) acnes*. [PLoS One. 13 \(1\): e0192051.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M290852:160715'

Printed on 04 May 2018