

Datasheet: MCA1076FT

Description:	MOUSE ANTI HUMAN CD62L:FITC
Specificity:	CD62L
Other names:	LECAM-1, L-SELECTIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	FMC46
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Bovine, Cynomolgus monkey, Rhesus Monkey, Dog N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid.		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	PHA stimulated lymphoblasts		

**External Database
Links**

UniProt:

[P14151](#) [Related reagents](#)

Entrez Gene:

[6402](#) SELL [Related reagents](#)

Synonyms

LNHR, LYAM1

Fusion Partners

Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line

Specificity

Mouse anti Human CD62L antibody, clone FMC46 recognizes human CD62L, also known as L-selectin, a 74-95 kDa member of the selectin family of adhesion receptors, which acts as a ligand for both CD62P (P-selectin) and CD62E (E-selectin). Human CD62L is constitutively expressed on most leucocytes including monocytes, granulocytes, lymphocytes, NK cells, bone marrow myeloid progenitor cells and on a subset of thymocytes.

CD62L plays an important role in leucocyte tethering and rolling on the endothelial cell surface and for the homing of naïve lymphocytes to lymph nodes and Peyer's patches via HEV. Neutrophils require a constant supply of this molecule on the cell surface for migration into peripheral tissues and adhesion to activated endothelium at sites of inflammation, where CD62L is rapidly shed as soluble L-selectin, but surface expression still remains.

The expression of CD62L is down regulated on lymphocytes and neutrophils by PMA stimulation.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul whole blood.

References

1. Pilarski, L.M. *et al.* (1991) FMC46, a cell protrusion-associated leukocyte adhesion molecule-1 epitope on human lymphocytes and thymocytes. [J Immunol. 147 \(1\): 136-43.](#)
2. Zola, H. *et al.* (1991) The expression of sub-population markers on B cells: a re-evaluation using high-sensitivity fluorescence flow cytometry. [Dis Markers. 9 \(2\): 103-18.](#)
3. Sopp, P. & Howard, C.J. (1997) Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. [Vet Immunol Immunopathol. 56 \(1-2\): 11-25.](#)
4. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) *Exp Anim.* 49 (2): 97-110.
5. Haanstra, K.G. *et al.* (2008) Characterization of naturally occurring CD4+CD25+ regulatory T cells in rhesus monkeys. [Transplantation 85:1185-92.](#)
6. Dalli, J. *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
7. Raposo, R.A. *et al.* (2011) Protein Kinase C and NF- κ B-Dependent CD4 Downregulation in Macrophages Induced by T Cell-Derived Soluble Factors: Consequences for HIV-1 Infection. [J Immunol. 187: 748-59.](#)
8. Hughes, S.F. *et al.* (2010) Total hip and knee replacement surgery results in changes in leukocyte and endothelial markers. [J Inflamm \(Lond\). 7:2.](#)
9. Bismarck, D. *et al.* (2012) Canine CD4+CD8+ double positive T cells in peripheral blood have features of activated T cells. [Vet Immunol Immunopathol. 149: 157-66.](#)
10. Hartley, A.N. & Tarleton, R.L. (2015) Chemokine receptor 7 (CCR7)-expression and IFN γ production define vaccine-specific canine T-cell subsets. [Vet Immunol Immunopathol. 164 \(3-4\): 127-36.](#)
11. Hayhoe, R.P. *et al.* (2006) Annexin 1 and its bioactive peptide inhibit neutrophil-endothelium interactions under flow: indication of distinct receptor involvement. [Blood. 107 \(5\): 2123-30.](#)

12. Urquhart, P. et al. (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther. 321 \(2\): 656-62.](#)
13. Aspinall, A.I. et al. (2010) CX(3)CR1 and vascular adhesion protein-1-dependent recruitment of CD16(+) monocytes across human liver sinusoidal endothelium. [Hepatology. 51 \(6\): 2030-9.](#)
14. Rothe, K. et al. (2017) Canine peripheral blood CD4⁺CD8⁺ double-positive Tcell subpopulations exhibit distinct Tcell phenotypes and effector functions. [Vet Immunol Immunopathol. 185: 48-56.](#)
15. Withers, S.S. et al. (2018) Multi-color flow cytometry for evaluating age-related changes in memory lymphocyte subsets in dogs. [Dev Comp Immunol. 87: 64-74.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M303531:170222'

Printed on 20 Jun 2018