

Datasheet: MCA1033GA

Description:	RAT ANTI MOUSE CD71
Specificity:	CD71
Other names:	TRANSFERRIN RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	YTA74.4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Concanavalin A activated mouse spleen cells.

**External Database
Links**

UniProt:

[Q62351](#) [Related reagents](#)

Entrez Gene:

[22042](#) Tfr [Related reagents](#)

Synonyms

Tfr

Fusion Partners

Spleen cells from an immunised DA rat were fused with cells of the Y3/Ag1.2.3 rat myeloma cell line.

Specificity

Rat anti Mouse CD71 antibody, clone YTA74.4 recognizes the mouse Transferrin receptor protein 1 also known as CD71 or TfR1. CD71 is a 763 amino acid glycoprotein homodimer of ~95 kDa subunits. CD71 is expressed by dividing cells, and functions as a transferrin receptor mediating uptake of iron.

Rat anti Mouse CD71 antibody, clone YTA74.4 blocks the binding of R17 217.1.3. and R17 208.2 anti-TFR monoclonal antibodies ([Trowbridge *et al.* 1982](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Lesley, J. *et al.* (1984) Expression of transferrin receptor on murine hematopoietic progenitors. [Cell Immunol. 83 \(1\): 14-25.](#)
2. Trowbridge, I.S. *et al.* (1982) Murine cell surface transferrin receptor: studies with an anti-receptor monoclonal antibody. [J Cell Physiol. 112 \(3\): 403-10.](#)
3. Millot, S. *et al.* (2010) Erythropoietin stimulates spleen BMP4-dependent stress erythropoiesis and partially corrects anemia in a mouse model of generalized inflammation. [Blood. 116: 6072-81.](#)
4. Kuo, Y.M. *et al.* (2004) Mislocalisation of hephaestin, a multicopper ferroxidase involved in basolateral intestinal iron transport, in the sex linked anaemia mouse. [Gut. 53: 201-6.](#)
5. Krysiak, K. *et al.* (2014) Reduced Levels of Hspa9 Attenuates Stat5 Activation in Mouse B-cells. [Exp Hematol. pii: S0301-472X\(14\)00817-0.](#)
6. Byun, M. *et al.* (2007) Cowpox virus exploits the endoplasmic reticulum retention pathway to inhibit MHC class I transport to the cell surface. [Cell Host Microbe. 2: 306-15.](#)
7. Ripich, T. and Jessberger, R. (2011) SWAP-70 regulates erythropoiesis by controlling $\alpha 4$ integrin. [Haematologica. 96: 1743-52.](#)
8. Niewoehner, J. *et al.* (2014) Increased brain penetration and potency of a therapeutic antibody using a monovalent molecular shuttle. [Neuron. 81: 49-60.](#)
9. Sands, S.A. *et al.* (2015) The habenula and iron metabolism in cerebral mouse models of multiple sclerosis. [Neurosci Lett. 606: 204-8.](#)
10. Baumann, B. *et al.* (2017) Conditional Müller Cell Ablation Leads to Retinal Iron Accumulation. [Invest Ophthalmol Vis Sci. 58 \(10\): 4223-34.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M315147:180503'

Printed on 05 May 2018