

Datasheet: MCA1148FT

Description:	MOUSE ANTI HUMAN CD71:FITC
Specificity:	CD71
Other names:	TRANSFERRIN RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	DF1513
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Rhesus Monkey, Mustelid, Ferret N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	KGI cell line.		

**External Database
Links**

UniProt:

[P02786](#) [Related reagents](#)

Entrez Gene:

[7037](#) TFRC [Related reagents](#)

Fusion Partners Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity **Mouse anti Human CD71 antibody, clone DF1513** recognizes the human CD71 cell surface antigen, a ~190 kDa homodimeric glycoprotein expressed by proliferating cells. CD71 is also known as the transferrin receptor. Mutation of the TFRC gene has been implicated in the development of Immunodeficiency 46 ([IMD46](#)) a combined immunodeficiency characterized by early onset chronic diarrhea, recurrent infections and intermittent neutropenia and thrombocytopenia ([Jabara *et al.* 2016](#))

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Sopper, S. *et al.* (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. [Cytometry. 29 \(4\): 351-62.](#)
2. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. [Vet Immunol Immunopathol. 132:109-15.](#)
3. Meng, J. *et al.* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. [PLoS One. 6\(3\):e17454.](#)
4. Stockwin, L.H. *et al.* (2009) Artemisinin dimer anticancer activity correlates with heme-catalyzed reactive oxygen species generation and endoplasmic reticulum stress induction. [Int J Cancer. 125: 1266-75.](#)
5. Janes, P.W. *et al.* (1999) Aggregation of lipid rafts accompanies signaling via the T cell antigen receptor. [J Cell Biol. 147: 447-61.](#)
6. Makoveichuk, E. *et al.* (2012) Inactivation of lipoprotein lipase occurs on the surface of THP-1 macrophages where oligomers of angiopoietin-like protein 4 are formed. [Biochem Biophys Res Commun. 425:138-43.](#)
7. Procaccini, C. *et al.* (2012) Leptin-induced mTOR activation defines a specific molecular and transcriptional signature controlling CD4+ effector T cell responses. [J Immunol. 189: 2941-53.](#)
8. Weissgerber, P. *et al.* (2003) Investigation of mechanisms involved in phagocytosis of Legionella pneumophila by human cells. [FEMS Microbiol Lett. 219 \(2\): 173-9.](#)
9. Trakarsanga, K. *et al.* (2017) An immortalized adult human erythroid line facilitates sustainable and scalable generation of functional red cells. [Nat Commun. 8: 14750.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Material Safety Datasheet documentation #10041 available at:

Information 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M304476:170406'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)