

## Datasheet: MCA1118

<b>Description:</b>	MOUSE ANTI HUMAN CD86
<b>Specificity:</b>	CD86
<b>Other names:</b>	B7-2
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	BU63
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/20 - 1/100
Immunohistology - Frozen	▪			1/20 - 1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human peripheral blood lymphocytes.

**External Database  
Links**

**UniProt:**

[P42081](#) [Related reagents](#)

**Entrez Gene:**

[942](#) CD86 [Related reagents](#)

---

**Synonyms**

CD28LG2

---

**Fusion Partners**

Spleen cells from immunised mice were fused with cells of the mouse P3.X63 Ag8653 myeloma cell line.

---

**Specificity**

**Mouse anti Human CD86 antibody, clone Bu63** recognizes human CD86 also known as B7-2, a type I transmembrane protein expressed by monocytes and activated B cells ([Engel et al. 1994](#)). CD86 acts as a co-stimulatory molecule along with CD80 ([Lanier et al. 1995](#)) and is a ligand for CD28 and CTLA-4 ([Azuma et al. 1993](#)).

CD86 is a member of the Immunoglobulin superfamily and carries an extracellular domain bearing both an [Ig-v-like](#) domain which contains the CTLA-4 binding site and an adjacent C2-like domain. CD86 plays an important role in co-stimulation of T cell proliferation ([Freeman et al. 1993](#)), IL-2 production ([Ribot et al. 2012](#)) and in the primary immune response ([Schultze et al. 1996](#)).

Domain depletion epitope mapping studies indicate that the binding site of Mouse anti Human CD86, [clone Bu63](#) is located within the Ig-v-like domain of human CD86 ([Jeanin et al. 1997](#)).

Studies suggest that CD86 along with CD80 may be exploited as receptors for adenovirus entry into cells ([Short et al. 2004](#) [2006](#)).

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

---

**Histology Positive  
Control Tissue**

Human Tonsil

---

**References**

1. McLellan, A.D. *et al.* (1999) Induction of dendritic cell costimulator molecule expression is suppressed by T cells in the absence of antigen-specific signalling: role of cluster formation, CD40 and HLA-class II for dendritic cell activation. [Immunology. 98 \(2\): 171-80.](#)
2. Nozawa, Y. *et al.* (1993) A novel monoclonal antibody (FUN-1) identifies an activation antigen in cells of the B-cell lineage and Reed-Sternberg cells. [J Pathol. 169 \(3\): 309-15.](#)
3. Goodyear, O. *et al.* (2010) Induction of a CD8+ T-cell response to the MAGE cancer testis antigen by combined treatment with azacitidine and sodium valproate in patients with acute myeloid leukemia and myelodysplasia. [Blood. 116: 1908-18.](#)
4. Angel, C.E. *et al.* (2006) Cutting edge: CD1a+ antigen-presenting cells in human dermis respond rapidly to CCR7 ligands. [J Immunol. 176 \(10\): 5730-4.](#)
5. Salte, T. *et al.* (2010) Increased intracellular growth of *Mycobacterium avium* in HIV-1 exposed monocyte-derived dendritic cells. [Microbes Infect. 13: 276-83.](#)
6. Adler, H.S. *et al.* (2010) Neuronal nitric oxide synthase modulates maturation of human dendritic cells. [J Immunol. 184: 6025-34.](#)
7. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. 12:2.](#)
8. Kapsogeorgou, E.K. *et al.* (2001) Functional expression of a costimulatory B7.2 (CD86) protein on human salivary gland epithelial cells that interacts with the CD28 receptor, but has reduced binding to CTLA4. [J Immunol. 166: 3107-13.](#)
9. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells

- facilitates T cell adhesion and migration. [J Immunol. 181: 6109-16.](#)
10. Urban, B.C. *et al.* (2001) A role for CD36 in the regulation of dendritic cell function. [Proc Natl Acad Sci U S A. 98: 8750-5.](#)
11. Zhan, H. *et al.* (2003) The immunomodulatory role of human conjunctival epithelial cells. [Invest Ophthalmol Vis Sci. 44: 3906-10.](#)
12. Sprater, F. *et al.* (2012) Expression of ESE-3 isoforms in immunogenic and tolerogenic human monocyte-derived dendritic cells. [PLoS One. 7 \(11\): e49577.](#)
13. McCarthy, N.E. *et al.* (2013) Proinflammatory V $\delta$ 2+ T Cells Populate the Human Intestinal Mucosa and Enhance IFN- $\gamma$  Production by Colonic  $\alpha\beta$  T Cells. [J Immunol. 191: 2752-63.](#)
14. Hofmann-Wellenhof, R. *et al.* (2004) Sunburn cell formation, dendritic cell migration, and immunomodulatory factor production after solar-simulated irradiation of sunscreen-treated human skin explants *in vitro*. [J Invest Dermatol. 123: 781-7.](#)
15. Rajkovic, I. *et al.* (2011) Differences in T-helper polarizing capability between human monocyte-derived dendritic cells and monocyte-derived Langerhans'-like cells. [Immunology. 132: 217-25.](#)

---

#### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

#### Shelf Life

18 months from date of despatch.

---

#### Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

#### Regulatory

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@549</a> , <a href="#">DyLight@649</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Human Anti Mouse IgG1 (HCA036...)	<a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

'M315174:180503'

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

**Printed on 05 May 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)