

Datasheet: MCA1118T

Description:	MOUSE ANTI HUMAN CD86
Specificity:	CD86
Other names:	B7-2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	BU63
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/20 - 1/100
Immunohistology - Frozen	■			1/20 - 1/100
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human peripheral blood lymphocytes.

**External Database
Links**

UniProt:

[P42081](#)

[Related reagents](#)

Entrez Gene:

[942](#)

CD86

[Related reagents](#)

Synonyms

CD28LG2

Fusion Partners

Spleen cells from immunised mice were fused with cells of the mouse P3.X63 Ag8653 myeloma cell line.

Specificity

Mouse anti Human CD86 antibody, clone Bu63 recognizes human CD86 also known as B7-2, a type I transmembrane protein expressed by monocytes and activated B cells ([Engel et al. 1994](#)). CD86 acts as a co-stimulatory molecule along with CD80 ([Lanier et al. 1995](#)) and is a ligand for CD28 and CTLA-4 ([Azuma et al. 1993](#)).

CD86 is a member of the Immunoglobulin superfamily and carries an extracellular domain bearing both an [Ig-v-like](#) domain which contains the CTLA-4 binding site and an adjacent C2-like domain. CD86 plays an important role in co-stimulation of T cell proliferation ([Freeman et al. 1993](#)), IL-2 production ([Ribot et al. 2012](#)) and in the primary immune response ([Schultze et al. 1996](#)).

Domain depletion epitope mapping studies indicate that the binding site of Mouse anti Human CD86, [clone Bu63](#) is located within the Ig-v-like domain of human CD86 ([Jeanin et al. 1997](#)).

Studies suggest that CD86 along with CD80 may be exploited as receptors for adenovirus entry into cells ([Short et al. 2004](#) [2006](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

**Histology Positive
Control Tissue**

Human Tonsil

References

1. McLellan, A.D. *et al.* (1999) Induction of dendritic cell costimulator molecule expression is suppressed by T cells in the absence of antigen-specific signalling: role of cluster formation, CD40 and HLA-class II for dendritic cell activation. [Immunology. 98 \(2\): 171-80.](#)
2. Nozawa, Y. *et al.* (1993) A novel monoclonal antibody (FUN-1) identifies an activation antigen in cells of the B-cell lineage and Reed-Sternberg cells. [J Pathol. 169 \(3\): 309-15.](#)
3. Goodyear, O. *et al.* (2010) Induction of a CD8+ T-cell response to the MAGE cancer testis antigen by combined treatment with azacitidine and sodium valproate in patients with acute myeloid leukemia and myelodysplasia. [Blood. 116: 1908-18.](#)
4. Angel, C.E. *et al.* (2006) Cutting edge: CD1a+ antigen-presenting cells in human dermis respond rapidly to CCR7 ligands. [J Immunol. 176 \(10\): 5730-4.](#)
5. Salte, T. *et al.* (2010) Increased intracellular growth of *Mycobacterium avium* in HIV-1 exposed monocyte-derived dendritic cells. [Microbes Infect. 13: 276-83.](#)
6. Adler, H.S. *et al.* (2010) Neuronal nitric oxide synthase modulates maturation of human dendritic cells. [J Immunol. 184: 6025-34.](#)
7. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. 12:2.](#)
8. Kapsogeorgou, E.K. *et al.* (2001) Functional expression of a costimulatory B7.2 (CD86) protein on human salivary gland epithelial cells that interacts with the CD28 receptor, but has reduced binding to CTLA4. [J Immunol. 166: 3107-13.](#)
9. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells

facilitates T cell adhesion and migration. [J Immunol. 181: 6109-16.](#)

10. Urban, B.C. *et al.* (2001) A role for CD36 in the regulation of dendritic cell function. [Proc Natl Acad Sci U S A. 98: 8750-5.](#)

11. Zhan, H. *et al.* (2003) The immunomodulatory role of human conjunctival epithelial cells. [Invest Ophthalmol Vis Sci. 44: 3906-10.](#)

12. Sprater, F. *et al.* (2012) Expression of ESE-3 isoforms in immunogenic and tolerogenic human monocyte-derived dendritic cells. [PLoS One. 7 \(11\): e49577.](#)

13. McCarthy, N.E. *et al.* (2013) Proinflammatory V δ 2+ T Cells Populate the Human Intestinal Mucosa and Enhance IFN- γ Production by Colonic $\alpha\beta$ T Cells. [J Immunol. 191: 2752-63.](#)

14. Hofmann-Wellenhof, R. *et al.* (2004) Sunburn cell formation, dendritic cell migration, and immunomodulatory factor production after solar-simulated irradiation of sunscreen-treated human skin explants *in vitro*. [J Invest Dermatol. 123: 781-7.](#)

15. Rajkovic, I. *et al.* (2011) Differences in T-helper polarizing capability between human monocyte-derived dendritic cells and monocyte-derived Langerhans'-like cells. [Immunology. 132: 217-25.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10057 available at: 10057: <https://www.bio-rad-antibodies.com/uploads/MSDS/10057.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

'M315175:180503'

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)