

Datasheet: MCA1270A488

Description:	MOUSE ANTI HUMAN CD13:Alexa Fluor® 488
Specificity:	CD13
Other names:	AMINOPEPTIDASE N
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	WM15
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human								
Species Cross Reactivity	Reacts with: Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species.								
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid								
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®488</td> <td>495</td> <td>519</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®488	495	519		
Fluorophore	Excitation Max (nm)	Emission Max (nm)							
Alexa Fluor®488	495	519							
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant								
Buffer Solution	Phosphate buffered saline								
Preservative	0.09% Sodium Azide								
Stabilisers	1% Bovine Serum Albumin								
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml								
Immunogen	Human AML cells.								

**External Database
Links**

UniProt:

[P15144](#) [Related reagents](#)

Entrez Gene:

[290](#) ANPEP [Related reagents](#)

Synonyms

APN, CD13, PEPN

Fusion Partners

Spleen cells from immunised BALB/c mice where fused with cells of the mouse NS1 myeloma cell line.

Specificity

Mouse anti Human CD13 antibody, clone WM15 recognizes human CD13 also known as aminopeptidase N. CD13 is a single pass type II glycosylated integral membrane protein with a predicted molecular mass of ~110 kDa and an apparent molecular mass of ~150 kDa expressed by granulocytes, monocytes, fibroblasts, endothelial cells and by myeloid leukaemia cells ([Bradstock et al. 1985](#)). CD13 acts as a major cell surface receptor for group 1 coronaviruses ([Breslin et al. 2003](#)) which bind to a critical sequence encompassing amino acid residues 288-295 ([Kolb et al. 1997](#)).

CD13 functions as an [aminopeptidase](#) enzyme, a metalloprotease present as both a membrane bound form and also a soluble aminopeptidase N.

Mouse anti Human CD13, clone WM15 inhibits infection of cells by human coronavirus ([Lachance et al. 1998](#)) but not hepatitis C virus ([Koutsoudakis et al. 2006](#)) and inhibits aminopeptidase N activity of the CD13 molecule ([Asmun et al. 1992](#))

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul whole blood

References

1. Bradstock, K.F. *et al.* (1985) Human myeloid differentiation antigens identified by monoclonal antibodies: expression on leukemic cells. [Pathology. 17 \(3\): 392-9.](#)
2. Bradstock, K.F. *et al.* (1985) Myeloid progenitor surface antigen identified by monoclonal antibody. [Br J Haematol. 61 \(1\): 11-20.](#)
3. Favaloro, E.J. *et al.* (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. [Br J Haematol. 69 \(2\): 163-71.](#)
4. Favaloro, E.J. (1991) CD-13 (gp150; aminopeptidase-N): co-expression on endothelial and haemopoietic cells with conservation of functional activity. [Immunol Cell Biol. 69 \(Pt 4\): 253-60.](#)
5. Favaloro, E.J. *et al.* (1993) The hepatobiliary disease marker serum alanine aminopeptidase predominantly comprises an isoform of the haematological myeloid differentiation antigen and leukaemia marker CD-13/gp150. [Clin Chim Acta. 220 \(1\): 81-90.](#)
6. Favaloro, E.J. *et al.* (1993) CD13 (GP150; aminopeptidase-N): predominant functional activity in blood is localized to plasma and is not cell-surface associated. [Exp Hematol. 21 \(13\): 1695-701.](#)
7. Tavoosidana, G. *et al.* (2011) Multiple recognition assay reveals prostasomes as promising plasma biomarkers for prostate cancer. [Proc Natl Acad Sci U S A. 108: 8809-14.](#)
8. Gredmark, S. *et al.* (2004) Human Cytomegalovirus Induces Inhibition of Macrophage Differentiation by Binding to Human Aminopeptidase N/CD13 [J Immunol. 173: 4897-907](#)
9. Grzywacz, B. *et al.* (2011) Natural killer-cell differentiation by myeloid progenitors. [Blood. 117: 3548-58.](#)
10. Stolzing, A. *et al.* (2008) Age-related changes in human bone marrow-derived mesenchymal stem cells: consequences for cell therapies. [Mech Ageing Dev. 129: 163-73.](#)
11. Silk, K.M. *et al.* (2012) Rapamycin conditioning of dendritic cells differentiated from human ES

- cells promotes a tolerogenic phenotype. [J Biomed Biotechnol. 2012:172420.](#)
12. Negussie, A.H. *et al.* (2010) Synthesis and in vitro evaluation of cyclic NGR peptide targeted thermally sensitive liposome. [J Control Release. 143: 265-73.](#)
13. Lassnig, C. *et al.* (2005) Development of a transgenic mouse model susceptible to human coronavirus 229E. [Proc Natl Acad Sci U S A. 102 \(23\): 8275-80.](#)
14. Thielitz, A. *et al.* (2004) Identification of extra- and intracellular alanyl aminopeptidases as new targets to modulate keratinocyte growth and differentiation. [Biochem Biophys Res Commun. 321 \(4\): 795-801.](#)
15. McCormack, E. *et al.* (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. [Blood. 121 \(7\): e34-42.](#)
16. Fiddler, C.A. *et al.* (2016) The Aminopeptidase CD13 Induces Homotypic Aggregation in Neutrophils and Impairs Collagen Invasion. [PLoS One. 11 \(7\): e0160108.](#)

Storage Store at +4°C for one month or at -20°C for longer.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 488 \(MCA928A488\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M299865:170105'

