

## Datasheet: MCA1271APC

<b>Description:</b>	MOUSE ANTI HUMAN CD33:APC
<b>Specificity:</b>	CD33
<b>Format:</b>	APC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	WM53
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human		
<b>Product Form</b>	Purified IgG conjugated to Allophycocyanin (APC) - lyophilised		
<b>Reconstitution</b>	Reconstitute with 1 ml distilled water		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	APC	650	661
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative</b>	0.09% Sodium Azide		
<b>Stabilisers</b>	1% Bovine Serum Albumin		
	5% Sucrose		
<b>Immunogen</b>	Human AML cells		
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P20138</a> <a href="#">Related reagents</a>		
	<b>Entrez Gene:</b>		

---

<b>Synonyms</b>	SIGLEC3
-----------------	---------

---

<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
------------------------	---

---

<b>Specificity</b>	<p><b>Mouse anti Human CD33 antibody, clone WM53</b> recognizes the human CD33 cell surface glycoprotein. This antigen, considered to be specific for the myeloid lineage, has also been reported to be present on cells of lymphoid origin.</p> <p>Mouse anti Human CD33 antibody, clone WM53 immunoprecipitates a protein of ~75kDa from myeloid cells, a smaller protein of approximately 67kDa has been observed in immunoprecipitates from lymphoid targets.</p>
--------------------	---

---

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
-----------------------	---

---

<b>References</b>	<ol style="list-style-type: none"><li>1. Favaloro, E.J. <i>et al.</i> (1987) Characterization of monoclonal antibodies to the human myeloid-differentiation antigen, gp67 (CD-33). <a href="#">Dis Markers. 5 (4): 215-25.</a></li><li>2. Favaloro, E.J. <i>et al.</i> (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. <a href="#">Br J Haematol. 69 (2): 163-71.</a></li><li>3. Hernández-Caselles, T. <i>et al.</i> (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. <a href="#">J Leukoc Biol. 79: 46-58.</a></li><li>4. Biedermann, B. <i>et al.</i> (2006) Analysis of the CD33-related siglec family reveals that Siglec-9 is an endocytic receptor expressed on subsets of acute myeloid leukemia cells and absent from normal hematopoietic progenitors. <a href="#">Leuk Res. 31: 211-20.</a></li><li>5. Lajaunias, F. <i>et al.</i> (2005) Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3-kinase-mediated intracellular signaling. <a href="#">Eur J Immunol. 35: 243-51.</a></li><li>6. Pietschmann, P. <i>et al.</i> (2000) Surface markers and transendothelial migration of dendritic cells from elderly subjects. <a href="#">Exp Gerontol. 35: 213-24.</a></li><li>7. Favaloro, E.J. <i>et al.</i> (1993) Differential expression of surface antigens on activated endothelium. <a href="#">Immunol Cell Biol. 71:571-81.</a></li><li>8. Yasukawa, T. <i>et al.</i> (2012) Simple detection of surface antigens on living cells by applying distinct cell positioning with negative dielectrophoresis. <a href="#">Anal Chem. 84 (20): 8830-6.</a></li><li>9. Hu, Z. <i>et al.</i> (2016) Self-assembled nanoparticles based on folic acid modified carboxymethyl chitosan conjugated with targeting antibody <a href="#">J Wuhan Univ of Technol-Mater. Sci. Ed. 31 (2): 446-53.</a></li><li>10. Dahl C <i>et al.</i> (2004) Human mast cells express receptors for IL-3, IL-5 and GM-CSF; a partial map of receptors on human mast cells cultured <i>in vitro</i>. <a href="#">Allergy. 59 (10): 1087-96.</a></li><li>11. Vamvakopoulos, J.E. &amp; Green, C. (2003) HMG-CoA reductase inhibition aborts functional differentiation and triggers apoptosis in cultured primary human monocytes: a potential mechanism of statin-mediated vasculoprotection. <a href="#">BMC Cardiovasc Disord. 3: 6.</a></li><li>12. Vamvakopoulos, J. <i>et al.</i> (2002) Genetic control of IL-1beta bioactivity through differential regulation of the IL-1 receptor antagonist. <a href="#">Eur J Immunol. 32 (10): 2988-96.</a></li><li>13. Lin, C.W. <i>et al.</i> (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. <a href="#">Blood. 106 (10): 3567-74.</a></li><li>14. McCormack E <i>et al.</i> (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. <a href="#">Blood. 121 (7): e34-42.</a></li></ol>
-------------------	--

---

**Storage** Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Shelf Life** 12 months from date of reconstitution.

---

**Health And Safety Information** Material Safety Datasheet documentation #10075 available at:  
10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:APC \(MCA928APC\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M299667:161221'

**Printed on 11 Jul 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)