

Datasheet: MCA1271GA

Description:	MOUSE ANTI HUMAN CD33
Specificity:	CD33
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	WM53
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human AML cells
External Database	UniProt:

Links

[P20138](#) [Related reagents](#)

Entrez Gene:

[945](#) CD33 [Related reagents](#)

Synonyms

SIGLEC3

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity

Mouse anti Human CD33 antibody, clone WM53 recognizes the human CD33 cell surface glycoprotein. This antigen, considered to be specific for the myeloid lineage, has also been reported to be present on cells of lymphoid origin.

Mouse anti Human CD33 antibody, clone WM53 immunoprecipitates a protein of ~75kDa from myeloid cells, a smaller protein of approximately 67kDa has been observed in immunoprecipitates from lymphoid targets.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells.

Immunohistology

* Overnight incubation recommended

Histology Positive Control Tissue

Lymphatic tissue

References

1. Favaloro, E.J. *et al.* (1987) Characterization of monoclonal antibodies to the human myeloid-differentiation antigen, gp67 (CD-33). [Dis Markers. 5 \(4\): 215-25.](#)
2. Favaloro, E.J. *et al.* (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. [Br J Haematol. 69 \(2\): 163-71.](#)
3. Hernández-Caselles, T. *et al.* (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. [J Leukoc Biol. 79: 46-58.](#)
4. Biedermann, B. *et al.* (2006) Analysis of the CD33-related siglec family reveals that Siglec-9 is an endocytic receptor expressed on subsets of acute myeloid leukemia cells and absent from normal hematopoietic progenitors. [Leuk Res. 31: 211-20.](#)
5. Lajaunias, F. *et al.* (2005) Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3-kinase-mediated intracellular signaling. [Eur J Immunol. 35: 243-51.](#)
6. Pietschmann, P. *et al.* (2000) Surface markers and transendothelial migration of dendritic cells from elderly subjects. [Exp Gerontol. 35: 213-24.](#)
7. Favaloro, E.J. *et al.* (1993) Differential expression of surface antigens on activated endothelium. [Immunol Cell Biol. 71:571-81.](#)
8. Yasukawa, T. *et al.* (2012) Simple detection of surface antigens on living cells by applying distinct cell positioning with negative dielectrophoresis. [Anal Chem. 84 \(20\): 8830-6.](#)
9. Hu, Z. *et al.* (2016) Self-assembled nanoparticles based on folic acid modified carboxymethyl chitosan conjugated with targeting antibody [J Wuhan Univ of Technol-Mater. Sci. Ed. 31 \(2\): 446-53.](#)
10. Dahl C *et al.* (2004) Human mast cells express receptors for IL-3, IL-5 and GM-CSF; a partial map of receptors on human mast cells cultured *in vitro*. [Allergy. 59 \(10\): 1087-96.](#)
11. Vamvakopoulos, J.E. & Green, C. (2003) HMG-CoA reductase inhibition aborts functional

differentiation and triggers apoptosis in cultured primary human monocytes: a potential mechanism of statin-mediated vasculoprotection. [BMC Cardiovasc Disord. 3: 6.](#)

12. Vamvakopoulos, J. *et al.* (2002) Genetic control of IL-1beta bioactivity through differential regulation of the IL-1 receptor antagonist. [Eur J Immunol. 32 \(10\): 2988-96.](#)

13. Lin, C.W. *et al.* (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. [Blood. 106 \(10\): 3567-74.](#)

14. McCormack E *et al.* (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. [Blood. 121 \(7\): e34-42.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Human Anti Mouse IgG1 (HCA036...) [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#),
[DyLight@649](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M317454:180702'

Printed on 11 Jul 2018
