

Datasheet: MCA1268PE

Description:	MOUSE ANTI HUMAN CD39:RPE
Specificity:	CD39
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	A1
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	PHA activated human lymphocytes		

External Database Links

UniProt:

[P49961](#)

[Related reagents](#)

Entrez Gene:

Synonyms	CD39
-----------------	------

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line
------------------------	--

Specificity	<p>Mouse anti Human CD39, clone A1 recognizes the human CD39 cell surface antigen, a ~70-100 kDa molecule expressed on peripheral blood B cells, T cells and monocytes, and weakly expressed by granulocytes.</p> <p>CD39 has intrinsic ecto-ATPase activity (Wang et al. 1996), and expression can be induced on T cells and increased on B cells, as a late activation antigen (Maliszewski et al. 1994).</p> <p>Mouse anti Human CD39, clone A1 has been shown to block MHC independent target cell recognition by hapten-specific CTL (Scholzen et al. 2009).</p>
--------------------	--

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
-----------------------	--

References	<ol style="list-style-type: none">1. Aversa, G.G. <i>et al.</i> (1988) Detection of a late lymphocyte activation marker by A1, a new monoclonal antibody. Transplant Proc. 20 (1): 49-52.2. Waugh, J.A. <i>et al.</i> (1989) Staining of normal and rejecting kidney using the activation panel. In: Leucocyte Typing IV. White cell differentiation antigens. Edited by Knapp, W. <i>et al.</i> Oxford University Press. p485.3. Aversa, G.G. and Hall, B.M. (1989) Activation panel antigen expression on PBL activated by PHA or in MLR. In: Leucocyte Typing IV. White cell differentiation antigens. Edited by Knapp, W. <i>et al.</i> Oxford University Press, p.498.4. Aversa, G.G. <i>et al.</i> (1989) Use of monoclonal antibodies to study in vivo and in vitro-activated lymphocytes. Transplant Proc. 21 (1 Pt 1): 349-50.5. Stein, H. <i>et al.</i> (1989) Activated Section report. In: Leucocyte Typing IV. White cell differentiation antigens. Edited by Knapp, W. <i>et al.</i> Oxford University Press, p.387.6. Suranyi, M.G. <i>et al.</i> (1991) Lymphocyte adhesion molecules in T cell-mediated lysis of human kidney cells. Kidney Int. 39 (2): 312-9.7. Stöckl, J. <i>et al.</i> (2001) Monomorphic molecules function as additional recognition structures on haptenated target cells for HLA-A1-restricted, hapten-specific CTL. J Immunol. 167 (5): 2724-33.8. Scholzen, A. <i>et al.</i> (2009) Plasmodium falciparum-mediated induction of human CD25Foxp3 CD4 T cells is independent of direct TCR stimulation and requires IL-2, IL-10 and TGFbeta. PLoS Pathog. 5: e1000543.9. Borsellino, G. <i>et al.</i> (2007) Expression of ectonucleotidase CD39 by Foxp3+ Treg cells: hydrolysis of extracellular ATP and immune suppression Blood. 110:1225-32.10. Mittag, D. <i>et al.</i> (2010) The effector T cell response to ryegrass pollen is counterregulated by simultaneous induction of regulatory T cells. J Immunol. 184: 4708-16.11. Loeuillet, C. <i>et al.</i> (2008) In vitro whole-genome analysis identifies a susceptibility locus for HIV-1. PLoS Biol. 6: e32.12. Rawstron, A.C. <i>et al.</i> (2010) Chronic lymphocytic leukaemia (CLL) and CLL-type monoclonal B-cell lymphocytosis (MBL) show differential expression of molecules involved in lymphoid tissue homing. Cytometry B Clin Cytom. 78 Suppl 1: S42-6.13. Alam, M.S. <i>et al.</i> (2009) CD73 is expressed by human regulatory T helper cells and suppresses proinflammatory cytokine production and Helicobacter felis-induced gastritis in mice. J Infect Dis. 199: 494-504.14. Moreno-Fernandez, M.E. <i>et al.</i> (2011) Regulatory T cells control HIV replication in activated T cells through a cAMP-dependent mechanism. Blood. 117: 5372-80.
-------------------	---

15. Guevara-Flores, A. *et al.* (2008) 5'-p-Fluorosulfonyl benzoyl adenosine inhibits an ecto-ATP-diphosphohydrolase in the tegument surface of *Taenia crassiceps* cysticerci. [Mol Biochem Parasitol. 162: 123-33.](#)
16. Glenn, J.R. *et al.* (2008) Raised levels of CD39 in leucocytosis result in marked inhibition of ADP-induced platelet aggregation via rapid ADP hydrolysis. [Platelets. 19: 59-69.](#)
17. Häusler SF *et al.* (2014) Anti-CD39 and anti-CD73 antibodies A1 and 7G2 improve targeted therapy in ovarian cancer by blocking adenosine-dependent immune evasion. [Am J Transl Res. 6 \(2\): 129-39.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

This product should be stored undiluted.

DO NOT FREEZE. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

12 months from date of reconstitution.

Health And Safety Information

Material Safety Datasheet documentation #10075 available at: 10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M306447:170612'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)