

Datasheet: MCA1260F

Description:	RAT ANTI MOUSE CD25:FITC
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	PC61.5.3
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	B6.1 CTL cell line.		
External Database Links	UniProt: P01590 Related reagents		

Entrez Gene:[16184](#) IL2ra [Related reagents](#)

Synonyms	IL2r
Fusion Partners	Spleen cells from immunized OFA rats were fused with cells of the P3X63Ag8.653 mouse myeloma cell line.
Specificity	Rat anti Mouse CD25 antibody, clone PC61.5.3 reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).
References	1. Moreau, J.L. <i>et al.</i> (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. Eur J Immunol. 17 (7): 929-35. 2. Hashimoto, N. <i>et al.</i> (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. Eur J Immunol. 16 (3): 317-20. 3. Lowenthal, J.W. <i>et al.</i> (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. J Immunol. 135 (6): 3988-94. 4. Ceredig, R. <i>et al.</i> (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. Nature 314: 98-100. 5. Yaqoob, P. & Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. Nutrition. 13 (7-8): 646-51. 6. Scotland, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. Blood. 118 (22): 5918-27. 7. Karali, D. <i>et al.</i> (2016) T cell regulation by <i>Phlomis lanata</i> protein extracts in mice. Pharm Biol. 54 (2): 207-14.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG1 NEGATIVE CONTROL:FITC \(MCA1211F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301459:170109'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)