

Datasheet: MCA1230A488

Description:	RAT ANTI MOUSE CD49d:Alexa Fluor® 488
Specificity:	CD49d
Other names:	INTEGRIN ALPHA 4 CHAIN, VLA-4
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	PS/2
Isotype:	IgG2b
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Species Cross Reactivity	Reacts with: Human N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®488	495	519
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	P815 DBA/2 murine mastocytoma cells.		

**External Database
Links**

UniProt:

[Q00651](#) [Related reagents](#)

Entrez Gene:

[16401](#) Itga4 [Related reagents](#)

Fusion Partners

Spleen cells from immunized Fisher rats were fused with SP2/0 mouse myeloma cells

Specificity

Rat anti Mouse CD49d monoclonal antibody, clone PS/2 recognizes murine alpha 4 integrin (CD49d), a ~150 kDa single pass type I membrane glycoprotein that can associate with either beta 1 integrin (CD29) or beta 7 integrin to form heterodimers CD49d/CD29 (VLA-4) and alpha4/beta7 (LPAM-1) respectively ([Holzmann et al. 1989](#)). CD49d is expressed on most lymphocytes, granulocytes, monocytes and thymocytes. The primary ligands for CD49d are CD106 (VCAM-1), fibronectin and MAdCAM-1 ([Sheppard et al. 1994](#)).

Clone PS/2 has also been reported to block the binding of CD49d to its ligands ([Andrew et al. 1994](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Miyake, K. *et al.* (1991) Evidence for a role of the integrin VLA-4 in lympho-hemopoiesis. [J Exp Med. 173 \(3\): 599-607.](#)
2. Miyake, K. *et al.* (1991) A VCAM-like adhesion molecule on murine bone marrow stromal cells mediates binding of lymphocyte precursors in culture. [J Cell Biol. 114 \(3\): 557-65.](#)
3. Andrew, D.P. *et al.* (1994) Distinct but overlapping epitopes are involved in alpha 4 beta 7-mediated adhesion to vascular cell adhesion molecule-1, mucosal addressin-1, fibronectin, and lymphocyte aggregation. [J Immunol. 153 \(9\): 3847-61.](#)
4. Bellingan, G.J. *et al.* (2002) Adhesion molecule-dependent mechanisms regulate the rate of macrophage clearance during the resolution of peritoneal inflammation. [J Exp Med. 196: 1515-21.](#)
5. Fukuoka, M. *et al.* (2000) Antiadhesive function of 130-kd glycoform of CD43 expressed in CD4 T-lymphocyte clones and transfectant cell lines. [Blood. 96: 4267-75.](#)
6. Hokibara, S. *et al.* (2000) Effects of monoclonal antibodies to adhesion molecules on eosinophilic myocarditis in Toxocara canis-infected CBA/J mice. [Clin Exp Immunol. 114: 236-44.](#)
7. Li, W. *et al.* (2008) Reduced alpha4beta1 integrin/VCAM-1 interactions lead to impaired pre-B cell repopulation in alpha 1,6-fucosyltransferase deficient mice. [Glycobiology. 18: 114-24.](#)
8. Liu, Z.J. *et al.* (1999) A novel role for H-Ras in the regulation of very late antigen-4 integrin and VCAM-1 via c-Myc-dependent and -independent mechanisms. [J Immunol. 163: 4901-8.](#)
9. Tchilian, E.Z. *et al.* (1997) Anti-alpha 4 integrin antibody induces apoptosis in murine thymocytes and staphylococcal enterotoxin B-activated lymph node T cells. [Immunology. 92: 321-7.](#)
10. Bowden, R.A. *et al.* (2002) Role of alpha4 integrin and VCAM-1 in CD18-independent neutrophil migration across mouse cardiac endothelium. [Circ Res. 90: 562-9.](#)
11. Hirata, T. *et al.* (2002) P-, E-, and L-selectin mediate migration of activated CD8+ T lymphocytes into inflamed skin. [J Immunol. 169: 4307-13.](#)
12. Vaz, R. *et al.* (2012) Fibronectin promotes migration, alignment and fusion in an in vitro myoblast cell model. [Cell Tissue Res. 348: 569-78.](#)
13. Gillberg, L. *et al.* (2013) Effective treatment of mouse experimental colitis by alpha 2 integrin antibody: comparison with alpha 4 antibody and conventional therapy. [Acta Physiol \(Oxf\). 207: 326-36.](#)
14. Eshghi, S. *et al.* (2007) Alpha4beta1 integrin and erythropoietin mediate temporally distinct

- steps in erythropoiesis: integrins in red cell development. [J Cell Biol. 177: 871-80.](#)
15. Maus, U.A. *et al.* (2004) Pneumolysin-induced lung injury is independent of leukocyte trafficking into the alveolar space. [J Immunol. 173: 1307-12.](#)
16. Enghofer, M. *et al.* (1998) Lymphocyte transfer in streptozotocin-induced diabetes: adhesion of donor cells to islet endothelium. [Am J Physiol. 274: E928-35.](#)
17. Zhang, Y. *et al.* (2012) Autotaxin through lysophosphatidic acid stimulates polarization, motility, and transendothelial migration of naive T cells. [J Immunol. 189: 3914-24.](#)
18. Ferrer, P. *et al.* (2005) Association between pterostilbene and quercetin inhibits metastatic activity of B16 melanoma. [Neoplasia. 7: 37-47.](#)
19. Tanneau, G.M. *et al.* (1999) Differential recruitment of T- and IgA B-lymphocytes in the developing mammary gland in relation to homing receptors and vascular addressins. [J Histochem Cytochem. 47: 1581-92.](#)
20. Omenetti, S. *et al.* (2015) Dysregulated intrahepatic CD4⁺ T-cell activation drives liver inflammation in ileitis-prone SAMP1/YitFc mice. [Cell Mol Gastroenterol Hepatol. 1 \(4\): 406-19.](#)
21. Chung, K.J. *et al.* (2017) A self-sustained loop of inflammation-driven inhibition of beige adipogenesis in obesity. [Nat Immunol. 18 \(6\): 654-664.](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Acknowledgements	The Alexa Fluor® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays, and are covered by pending and issued patents.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M289297:170105'

Printed on 01 Aug 2018