

Datasheet: MCA1209PE

Description:	MOUSE IgG1 NEGATIVE CONTROL:RPE
Specificity:	MOUSE IgG1 NEGATIVE CONTROL
Format:	RPE
Product Type:	Negative/Isotype Control
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.

Target Species	Negative Control		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	Human T lymphocytes.		
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.		
Specificity	Mouse IgG1 Negative Control antibody is suitable for use as a negative control to assess non-specific binding of mouse IgG1 antibodies to target cells. Mouse IgG1 Negative Control antibody has been tested and found to be negative on the following rat cell types, peripheral blood		

leucocytes, thymocytes, splenocytes and macrophages.

Clone F8-11-13 recognises the human CD45RA antigen, and therefore human leucocytes may be used as a positive control for this product. NOT SUITABLE FOR USE AS A NEGATIVE CONTROL ON HUMAN TISSUES

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
-----------------------	---

References	1. Weiss, D.J. <i>et al.</i> (2008) Bovine monocyte TLR2 receptors differentially regulate the intracellular fate of <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> and <i>Mycobacterium avium</i> subsp. <i>avium</i>. J Leukoc Biol. 83 (1): 48-55. 2. Chen, W. <i>et al.</i> (2009) Expression of toll-like receptor 4 in uvea-resident tissue macrophages during endotoxin-induced uveitis. Mol Vis. 15: 619-28. 3. Safeukui I <i>et al.</i> (2015) Malaria induces anemia through CD8+ T cell-dependent parasite clearance and erythrocyte removal in the spleen. MBio. 6 (1) pii: e02493-14. 4. Aricha, R. <i>et al.</i> (2016) Suppression of experimental autoimmune myasthenia gravis by autologous T regulatory cells. J Autoimmun. 67: 57-64. 5. Wattegedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. Vet Res. 48 (1): 20.
-------------------	---

Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	---

Shelf Life	6 months from date of reconstitution.
-------------------	---------------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301163:170106'

Printed on 21 Jun 2018