

Datasheet: MCA1210PE

Description:	MOUSE IgG2a NEGATIVE CONTROL:RPE
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	RPE
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

* It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.

Target Species	Negative Control		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by ion exchange chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	Human T lymphocytes.		
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of NS1 mouse myeloma cell line.		
Specificity	Mouse IgG2a Negative Control is suitable for use as a matched isotype negative control reagent to assess the level of non-specific binding of mouse IgG2a monoclonal antibodies to rat cells.		

antibody has been tested and found to be negative on the following rat cells – peripheral blood leucocytes, thymocytes, splenocytes and macrophages.

MCA1210 recognizes the human CD45 antigen and therefore human leucocytes may be used as a positive control for this product.

THIS PRODUCT IS NOT SUITABLE FOR USE AS A NEGATIVE CONTROL ON HUMAN TISSUES.

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Fijak, M. <i>et al.</i> (2015) Influence of Testosterone on Inflammatory Response in Testicular Cells and Expression of Transcription Factor Foxp3 in T Cells. Am J Reprod Immunol. 74 (1): 12-25.
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:RPE \(MCA929PE\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M301165:170106'

Printed on 21 Jun 2018