

Datasheet: MCA1211B

Description:	RAT IgG1 NEGATIVE CONTROL:Biotin		
Specificity:	RAT IgG1 NEGATIVE CONTROL		
Format:	Biotin		
Product Type:	Negative/Isotype Control		
Isotype:	lgG1		
Quantity:	0.1 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

* It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.

Target Species	Negative Control	
Product Form	Purified IgG conjugated to Biotin - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A	
Buffer Solution	Phosphate buffered saline	
Preservative	0.09% Sodium Azide	
Stabilisers	1% Bovine Serum Albumin	
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml	
Fusion Partners	Spleen cells from immunized PVG-RT1u rat were fused with cells of cell line.	the rat Y3/Ag1.2.3 myeloma
Specificity	Rat IgG1 Negative Control antibody is suitable for the measurement monoclonal antibodies of isotype IgG1 to target cells. Rat IgG1 Negatintendend for use on mouse tissues.	· · · · · · · · · · · · · · · · · · ·
	N.B. This Rat IgG1 Negative Control antibody recognizes a human of therefore cannot be used as a negative control in this species.	cell surface marker, and

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.			
References	 Grimm, M. et al. (2010) Evaluation of immunological escape mechanisms in a mouse model of colorectal liver metastases. <u>BMC Cancer. 10:82.</u> Tighe, R.M. et al. (2011) Ozone Inhalation Promotes CX3CR1-Dependent Maturation of Resident Lung Macrophages That Limit Oxidative Stress and Inflammation. <u>J Immunol. 187: 4800-8.</u> Heidt, T. et al. (2016) Molecular Imaging of Activated Platelets Allows the Detection of Pulmonary Embolism with Magnetic Resonance Imaging. <u>Sci Rep. 6: 25044.</u> 			
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted.			
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.			
Shelf Life	18 months from date of despatch.			
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf			
Regulatory	For research purposes only			

Related Products

Recommended Negative Controls

RAT IgG1 NEGATIVE CONTROL:BIOTIN (MCA6004B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M258015:140617'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint