

# Datasheet: MCA1212D649

Description:	RAT IgG2a NEGATIVE CONTROL:DyLight® 649
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	DyLight®649
Product Type:	Negative/Isotype Control
Isotype:	lgG2a
Quantity:	0.1 mg

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	NO	Not Determined	Suggested Dilution
Flow Cytometry	•			*
Where this antibody has no	t been tes	sted for use	e in a particular technique	e this does not necessarily

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.\* It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test reagents.

Target Species	Negative Control				
Product Form	Purified IgG conjugated to DyLight <sup>®</sup> 649 - liquid				
Max Ex/Em	Fluorophore Excitation Max (nm) Emission Max (nm)				
	Dylight®649 654 673				
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide				
Stabilisers	1% Bovine Serum Albumin				
Approx. Protein Concentrations	IgG concentration 0.1mg/ml				
Immunogen	Human lymphocytes.				
Fusion Partners	Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line.				
Specificity	Rat IgG2a Negative Control antibody is suitable for the assessment of the level of non-specific binding of rat IgG2a monoclonal antibodies to mouse cells.				

Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative

control with dog cells.

N.B. This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines.

#### References

- 1. Sumagin, R. et al. (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. Am J Physiol Heart Circ Physiol. 295: H969-H977.
- 2. Chiu, W.C. et al. (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. J Nutr Biochem. 20: 254-60.
- 3. Guilloteau, L.A. et al. (2003) Nramp1 is not a major determinant in the control of Brucella melitensis infection in mice. Infect Immun. 71: 621-8.
- 4. Stapleton, T.W. et al. (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. Tissue Eng Part A. 17: 231-42.
- 5. Park, S.W. et al. (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. Am J Physiol Renal Physiol. 303: F721-32.
- 6. Schmidt, E.P. et al. (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. Nat Med. 18 (8): 1217-23.
- 7. McConnell, M.J. et al. (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. Proc Natl Acad Sci U S A. 106: 6784-9.
- 8. Rabadi MM et al. (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. Am J Physiol Renal Physiol. Jun 22: ajprenal.00254.2016. [Epub ahead of print]

### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
Acknowledgements	DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>
Regulatory	For research purposes only

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

'M299070:161213'

Printed on 20 Jun 2018