

Datasheet: MCA1360A647

Description:	MOUSE ANTI V5-TAG:Alexa Fluor® 647
Specificity:	V5-TAG
Other names:	PK-TAG
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	SV5-Pk1
Isotype:	IgG2a
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Viral		
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	Paramyxovirus Simian-Virus 5 (SV5)		
External Database Links	UniProt: P11207 Related reagents		

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the SP2/0 Ag14 myeloma cell line.
Specificity	Mouse anti V5-Tag, clone SV5-Pk1 recognizes the sequence, IPNPLLGL, present on the P/V proteins of the paramyxovirus, SV5 (Dunn et al.1999). Clone SV5-Pk1 is used to detect recombinant proteins, some of which include transmembrane and secreted proteins, that have labelled with tags containing this sequence (Randall et al.1993 and Zhao et al. 2005).
References	1. Southern, J.A. <i>et al.</i> (1991) Identification of an epitope on the P and V proteins of simian virus 5 that distinguishes between two isolates with different biological characteristics. J Gen Virol. 72 (Pt 7): 1551-7. 2. Orime, K. <i>et al.</i> (2013) Trefoil Factor 2 Promotes Cell Proliferation in Pancreatic β-Cells through CXCR-4-Mediated ERK1/2 Phosphorylation. Endocrinology. 154: 54-64. 3. Randall, R.E. <i>et al.</i> (1993) Two-tag purification of recombinant proteins for the construction of solid matrix-antibody-antigen (SMAA) complexes as vaccines. Vaccine. 11 (12): 1247-52. 4. Randall, R.E. <i>et al.</i> (1994) Purification of antibody-antigen complexes containing recombinant SIV proteins: comparison of antigen and antibody-antigen complexes for immune priming. Vaccine. 12 (4): 351-8. 5. Hanke, T. <i>et al.</i> (1995) Attachment of an oligopeptide epitope to the C-terminus of recombinant SIV gp160 facilitates the construction of SMAA complexes while preserving CD4 binding. J Virol Methods. 53 (1): 149-56. 6. Jaffray, E. <i>et al.</i> (1995) Domain organization of I kappa B alpha and sites of interaction with NF-kappa B p65. Mol Cell Biol. 15 (4): 2166-72. 7. Rodriguez, M.S. <i>et al.</i> (1995) Inducible degradation of I kappa B alpha in vitro and in vivo requires the acidic C-terminal domain of the protein. Mol Cell Biol. 15 (5): 2413-9. 8. Arenzana-Seisdedos, F. <i>et al.</i> (1995) Inducible nuclear expression of newly synthesized I kappa B alpha negatively regulates DNA-binding and transcriptional activities of NF-kappa B. Mol Cell Biol. 15 (5): 2689-96. 9. Hirst, K. <i>et al.</i> (1994) The transcription factor, the Cdk, its cyclin and their regulator: directing the transcriptional response to a nutritional signal. EMBO J. 13 (22): 5410-20. 10. Dunn, C. <i>et al.</i> (1999) Fine mapping of the binding sites of monoclonal antibodies raised against the Pk tag. J Immunol Methods. 224 (1-2): 141-50. 11. Young, D.F. <i>et al.</i> (2001) Single amino acid substitution in the V protein of simian virus 5 differentiates its ability to block interferon signaling in human and murine cells. J Virol. 75 (7): 3363-70. 12. Sanchez Garcia, J. <i>et al.</i> (2004) The C-terminal zinc finger of the catalytic subunit of DNA polymerase delta is responsible for direct interaction with the B-subunit. Nucleic Acids Res. 32 (10): 3005-16. 13. Herskowitz, J.H. <i>et al.</i> (2011) Rho kinase II phosphorylation of the lipoprotein receptor LR11/SORLA alters amyloid-beta production. J Biol Chem. 286 (8): 6117-27. 14. Liebau, M.C. <i>et al.</i> (2011) Nephrocystin-4 regulates Pyk2-induced tyrosine phosphorylation of Nephrocystin-1 to control targeting to monocilia. J Biol Chem. 286: 14237-45. 15. Björk, J.K. <i>et al.</i> (2010) miR-18, a member of Oncomir-1, targets heat shock transcription factor 2 in spermatogenesis. Development. 137(19):3177-84. 16. Boggio, R. <i>et al.</i> (2007) Targeting SUMO E1 to ubiquitin ligases: a viral strategy to counteract sumoylation. J Biol Chem. 282: 15376-82. 17. Gallazzini, M. <i>et al.</i> (2011) High NaCl-induced activation of CDK5 increases phosphorylation of the osmoprotective transcription factor TonEBP/OREBP at threonine 135, which contributes to its rapid nuclear localization. Mol Biol Cell. 22: 703-14. 18. Hadler, K.S. <i>et al.</i> (2008) Identification of a non-purple tartrate-resistant acid phosphatase: an evolutionary link to Ser/Thr protein phosphatases? BMC Res Notes. 1: 78. 19. Zhao, A. <i>et al.</i> (2011) Rapid isolation of high-affinity human antibodies against the tumor vascular marker Endosialin/TEM1, using a paired yeast-display/secretory scFv library platform. J

[Immunol Methods. 363: 221-32.](#)

20. Patino, G.A. *et al.* (2011) Voltage-Gated Na⁺ Channel β 1B: A Secreted Cell Adhesion Molecule Involved in Human Epilepsy. [J Neurosci. 31: 14577-91.](#)
21. Gatherer, D. *et al.* (2011) High-resolution human cytomegalovirus transcriptome. [Proc Natl Acad Sci U S A. 108: 19755-60.](#)
22. Mahuzier, A. *et al.* (2012) Dishevelled stabilization by the ciliopathy protein Rpgrip1l is essential for planar cell polarity. [J Cell Biol. 198: 927-40.](#)
23. Zhao, C. *et al.* (2005) Human ISG15 conjugation targets both IFN-induced and constitutively expressed proteins functioning in diverse cellular pathways. [Proc Natl Acad Sci U S A. 102:10200-5](#)
24. Singh, A. *et al.* (2014) Trypanosome MKT1 and the RNA-binding protein ZC3H11: interactions and potential roles in post-transcriptional regulatory networks. [Nucleic Acids Res. 42: 4652-68.](#)
25. Mui, M.Z. *et al.* (2015) The Human Adenovirus Type 5 E4orf4 Protein Targets Two Phosphatase Regulators of the Hippo Signaling Pathway. [J Virol. 89 \(17\): 8855-70.](#)
26. Shi X *et al.* (2016) Bunyamwera orthobunyavirus glycoprotein precursor is processed by cellular signal peptidase and signal peptide peptidase. [Proc Natl Acad Sci U S A. 113 \(31\): 8825-30.](#)
27. Ng, M.Y. *et al.* (2017) Activation of MAPK/ERK signaling by *Burkholderia pseudomallei* cycle inhibiting factor (Cif). [PLoS One. 12 \(2\): e0171464.](#)
28. Voskarides, K. *et al.* (2017) A functional variant in NEPH3 gene confers high risk of renal failure in primary hematuric glomerulopathies. Evidence for predisposition to microalbuminuria in the general population. [PLoS One. 12 \(3\): e0174274.](#)
29. Malik, S. *et al.* (2015) Adrenocorticotrophic Hormone (ACTH) Responses Require Actions of the Melanocortin-2 Receptor Accessory Protein on the Extracellular Surface of the Plasma Membrane. [J Biol Chem. 290 \(46\): 27972-85.](#)
30. Carrocci, T.J. *et al.* (2017) SF3b1 mutations associated with myelodysplastic syndromes alter the fidelity of branchsite selection in yeast. [Nucleic Acids Res. 45 \(8\): 4837-4852.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is manufactured under an exclusive license from the University of St. Andrews, UK.

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

RegulatoryFor research purposes only

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M300567:170105'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)