

Datasheet: MCA1450F

Description:	MOUSE ANTI HUMAN CD119:FITC
Specificity:	CD119
Other names:	IFN GAMMA RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	BB1E2
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Bovine N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by ion exchange chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Fusion protein of the extracellular part of the human interferon-gamma receptor (AA: Ala17-Gly245) conjugated to 11 AA of the c-myc proto-oncogene (for affinity purification).		

**External Database
Links**

UniProt:

[P15260](#)

[Related reagents](#)

Entrez Gene:

[3459](#)

IFNGR1

[Related reagents](#)

Fusion Partners

Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0-Ag14 myeloma cell line.

Specificity

Mouse anti Human CD119 antibody, clone BB1E2 recognizes the human gamma-interferon receptor, also known as CD119. CD119 is expressed by monocytes, and very weakly by resting peripheral blood lymphocytes.

Mouse anti Human CD119 antibody, clone BB1E2 recognizes a single protein band of ~43 kDa in western blotting under non-reducing conditions against protein comprising the glycosylated extracellular domain of the human interferon-gamma receptor fused to c-myc, and a MW of 27kDa against a recombinant protein corresponding to the non-glycosylated domain fused to c-myc.

Mouse anti Human CD119 antibody, clone BB1E2 does not neutralize biological activity of interferon gamma.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Entrican G *et al.* (2002) Gamma interferon fails to induce expression of indoleamine 2,3-dioxygenase and does not control the growth of *Chlamydomyphila abortus* in BeWo trophoblast cells. [Infect Immun. 70 \(5\): 2690-3.](#)
2. Sopp, P. *et al.* (2007) Cross-reactivity of mAbs to human CD antigens with cells from cattle. [Vet Immunol Immunopathol. 119: 106-14.](#)
3. López-Herrera, G. *et al.* (2004) Severe combined immunodeficiency syndrome associated with colonic stenosis. [Arch Med Res. 35: 348-58.](#)
4. Garibay-Escobar, A. *et al.* (2003) Integrated measurements by flow cytometry of the cytokines IL-2, IFN-gamma, IL-12, TNF-alpha and functional evaluation of their receptors in human blood. [J Immunol Methods. 280: 73-88.](#)
5. Jankovicova, K. *et al.* (2012) Interferon gamma receptor expression on granulocytes of cardiac surgical patients is modulated differently by the type of cardiopulmonary bypass used. [Perfusion. 27: 49-55.](#)
6. Diaz-Romero, J. *et al.* (2005) Immunophenotypic analysis of human articular chondrocytes: changes in surface markers associated with cell expansion in monolayer culture. [J Cell Physiol. 202: 731-42.](#)
7. Garibay-Escobar, A. *et al.* (2003) Integrated measurements by flow cytometry of the cytokines IL-2, IFN-gamma, IL-12, TNF-alpha and functional evaluation of their receptors in human blood. [J Immunol Methods. 280: 73-88.](#)
8. Ghaemmaghami, A.M. *et al.* (2002) The proteolytic activity of the major dust mite allergen Der p 1 conditions dendritic cells to produce less interleukin-12: allergen-induced Th2 bias determined at the dendritic cell level. [Clin Exp Allergy. 32: 1468-75.](#)
9. Yim, J.H. *et al.* (2003) The role of interferon regulatory factor-1 and interferon regulatory factor-2 in IFN-gamma growth inhibition of human breast carcinoma cell lines. [J Interferon Cytokine Res. 23: 501-11.](#)

Further Reading

1. Michiels, L. *et al.* (1998) The soluble extracellular portion of the human interferon-gamma receptor is a valid substitute for evaluating binding characteristics and for neutralizing the biological activity of this cytokine. [Int J Biochem Cell Biol. 30: 505-16.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:FITC \(MCA929F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M301251:170109'

Printed on 15 May 2018