

## Datasheet: MCA1439PE

<b>Description:</b>	RAT ANTI MOUSE CD19:RPE
<b>Specificity:</b>	CD19
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	6D5
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	100 TESTS

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse		
<b>Product Form</b>	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
<b>Reconstitution</b>	Reconstitute with 1 ml distilled water		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	RPE 488nm laser	496	578
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative</b>	0.09% Sodium Azide		
<b>Stabilisers</b>	1% Bovine Serum Albumin		
	5% Sucrose		
<b>Immunogen</b>	Human K562 cell line transfected with murine CD19.		
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P25918</a> <a href="#">Related reagents</a>		
	<b>Entrez Gene:</b>		

---

<b>Fusion Partners</b>	Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line.
------------------------	--

---

<b>Specificity</b>	<b>Rat anti Mouse CD19 antibody, clone 6D5</b> recognizes the murine CD19 cell surface antigen, a 95kDa glycoprotein expressed by B lymphocytes. Clone 6D5 has been shown to recognize the same epitope as clone ID3 in cross-competition assays.
--------------------	---

---

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.  The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ( <a href="#">BUF041A/B</a> ).
-----------------------	--

---

<b>References</b>	<ol style="list-style-type: none"><li>1. Vernooy, J.H. <i>et al.</i> (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. <a href="#">Am J Respir Cell Mol Biol. 26 (1): 152-9.</a></li><li>2. Andrew, D. and Aspinall. R. (2001) Il-7 and not stem cell factor reverses both the increase in apoptosis and the decline in thymopoiesis seen in aged mice. <a href="#">J Immunol. 166: 1524-30.</a></li><li>3. Bermudez-Fajardo, A. <i>et al.</i> (2011) The effect of Chlamydomphila pneumoniae Major Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune responses. <a href="#">Immunobiology. 216: 152-63.</a></li><li>4. De Jesus, M. <i>et al.</i> (2009) Galactoxylomannan-mediated immunological paralysis results from specific B cell depletion in the context of widespread immune system damage. <a href="#">J Immunol. 183: 3885-94.</a></li><li>5. Jégou, J.F. <i>et al.</i> (2007) C3d binding to the myelin oligodendrocyte glycoprotein results in an exacerbated experimental autoimmune encephalomyelitis. <a href="#">J Immunol. 178: 3323-31.</a></li><li>6. Starck, J. <i>et al.</i> (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. <a href="#">Blood. 116: 4795-805.</a></li><li>7. Scotland, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <a href="#">Blood. 118 (22): 5918-27.</a></li><li>8. White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. <a href="#">J Immunol. 188: 287-93.</a></li><li>9. Reynaud, J.M. <i>et al.</i> (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via Toll-like receptor 9. <a href="#">J Virol. 88: 5421-36.</a></li><li>10. Candolfi, M. <i>et al.</i> (2011) B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. <a href="#">Neoplasia. 13: 947-60.</a></li><li>11. Takabayashi, H. <i>et al.</i> (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of mice. <a href="#">Gastroenterology. 147: 396-406.e7.</a></li><li>12. Weiss-Gayet, M. <i>et al.</i> (2016) Notch Stimulates Both Self-Renewal and Lineage Plasticity in a Subset of Murine CD9High Committed Megakaryocytic Progenitors. <a href="#">PLoS One. 11 (4): e0153860.</a></li><li>13. Meng, Q.H. &amp; White, H.N. (2017) CD21<sup>int</sup> CD23<sup>+</sup> follicular B cells express antigen-specific secretory IgM mRNA as primary and memory responses. <a href="#">Immunology. 151 (2): 211-8.</a></li><li>14. Baer, A. <i>et al.</i> (2016) Protein Phosphatase-1 regulates Rift Valley fever virus replication. <a href="#">Antiviral Res. 127: 79-89.</a></li><li>15. Mccubbrey, A.L. <i>et al.</i> (2016) MicroRNA-34a Negatively Regulates Efferocytosis by Tissue Macrophages in Part via SIRT1. <a href="#">J Immunol. 196 (3): 1366-75.</a></li></ol>
-------------------	--

---

<b>Storage</b>	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.
----------------	--

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Shelf Life</b>	12 months from date of reconstitution.
-------------------	--

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10075 available at: 10075: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA1212PE\)](#)

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M300696:170106'

**Printed on 01 Aug 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)