

Datasheet: MCA1578F

Description:	MOUSE ANTI HUMAN CD34:FITC
Specificity:	CD34 CLASS III
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	581
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human								
Species Cross Reactivity	Reacts with: Cynomolgus monkey N.B. Antibody reactivity and working conditions may vary between species.								
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid.								
Max Ex/Em	<table><tr><th>Fluorophore</th><th>Excitation Max (nm)</th><th>Emission Max (nm)</th></tr><tr><td>FITC</td><td>490</td><td>525</td></tr></table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525		
Fluorophore	Excitation Max (nm)	Emission Max (nm)							
FITC	490	525							
Preparation	Purified IgG prepared by affinity chromatography on Protein A.								
Buffer Solution	Phosphate buffered saline								
Preservative	0.09% Sodium Azide								
Stabilisers	0.2% Bovine Serum Albumin								
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml.								
External Database Links	UniProt: P28906 Related reagents								

Entrez Gene:

[947](#) CD34 [Related reagents](#)

Specificity	Mouse anti Human CD34 antibody, clone 581 recognizes the human CD34 cell surface antigen also known as Hematopoietic progenitor cell antigen CD34. Clone 581 binds to an epitope of human CD34 resistant to neuraminadase, chymopapain and glycoprotease treatments. The binding epitopes are classified according to their resistance to enzymatic degradation and antibodies assigned according to this and cross inhibition studies. Clone 581 is classified as class III (Lanza et al. 1999). CD34 is a 385 amino acid pro-peptide with the 31 amino acid signal peptide cleaved to yield the mature form of CD34. CD34 is a single pass type I ~105-120 kD highly glycosylated transmembrane glycoprotein, expressed by hematopoietic stem cells and small vessel endothelial cells. CD34 contains a number of sialylated residues and has a mucin-like structure (Simmons et al. 1992). A truncated form of the CD34 molecule has also been identified lacking the carboxy-terminal region of the intracellular sequence, also bearing a sequence switch in the remaining cytoplasmic region (Nakamura et al. 1993). Clone 581 is expected to react with both isoforms of the CD34 molecule as it recognizes an epitope located in the extracellular region of human CD34. Mouse anti Human CD34 antibody, clone 581 has been used successfully for the immunohistochemical detection of stem-like cells present in hepatoblastoma lesions on cryostat sections (Fiegel et al. 2004).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Civin, C.I. <i>et al.</i> (1984) Antigenic analysis of hematopoiesis. III. A hematopoietic progenitor cell surface antigen defined by a monoclonal antibody raised against KG-1a cells. J Immunol. 133 (1): 157-65. 2. Yoshino, N. <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. Exp. Anim. 49: 97-110. 3. Maumus, M. <i>et al.</i> (2011) Native human adipose stromal cells: localization, morphology and phenotype. Int J Obes (Lond). 35 (9): 1141-53. 4. Fiegel, H.C. <i>et al.</i> (2004) Stem-like cells in human hepatoblastoma. J Histochem Cytochem. 52 (11): 1495-501.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301552:170109'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)