

Datasheet: MCA1511PE

Description:	RAT ANTI MOUSE CD38:RPE
Specificity:	CD38
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	90
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Buffer Solution	Phosphate buffered saline		
Preservative	0.1% Sodium Azide		
Stabilisers	0.5% Bovine Serum Albumin		
Immunogen	Mouse pre-B cells derived from IL-7 dependent bone marrow cultures.		
External Database Links	UniProt: P56528 Related reagents		
	Entrez Gene: 12494 Cd38 Related reagents		
Fusion Partners	Lymph node cells from an immunized rat were fused with cells of the m myeloma cell line.		

Specificity	Rat anti Mouse CD38 antibody, clone 90 recognizes the murine CD38 cell surface antigen, a ~42-46kDa glycoprotein expressed by B lymphocytes, a proportion of peripheral T lymphocytes and by peritoneal macrophages. In the mouse CD38 is expressed by follicular B cells, is down regulated on germinal centre B cells, and not expressed by mature plasma cells. This pattern of expression contrasts strikingly with that seen in the human.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).
References	1. Lund, F. <i>et al.</i> (1995) Murine CD38: an immunoregulatory ectoenzyme. Immunol. Today. 16: 469. 2. Oliver, A.M. <i>et al.</i> (1997) Mouse CD38 is down-regulated on germinal center B cells and mature plasma cells. J. Immunol. 158: 1108-1115.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	6 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10304 available at: 10304: https://www.bio-rad-antibodies.com/uploads/MSDS/10304.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA1212PE\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M304582:170410'

Printed on 02 May 2018