

Datasheet: MCA1615

Description:	MOUSE ANTI HUMAN CD54
Specificity:	CD54
Other names:	ICAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	15.2
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/50 - 1/100
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Pig N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	Human monocytes
External Database Links	<p>UniProt: P05362 Related reagents</p> <p>Entrez Gene: 3383 ICAM1 Related reagents</p>
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/O-Ag14 myeloma cell line
Specificity	<p>Mouse anti Human CD54 antibody, clone 15.2 recognizes the human CD54 cell surface antigen also known as intracellular Adhesion Molecule -1 (ICAM-1) or Major group rhinovirus receptor.</p> <p>CD54 is expressed by many cells following activation by inflammatory mediators. It is a 505 amino acid with an additional 27 amino acid signal peptide ~90 kDa single pass type I transmembrane glycoprotein bearing 5 Ig-like C2-type domains.</p> <p>Mouse anti Human CD54 antibody, clone 15.2 blocks CD54 function (Berendt <i>et al.</i> 1992). Mouse anti Human CD54 antibody, clone 15.2 binds to an epitope on the N-terminal Ig-like domain within a region designated the L43 loop (Chakravorty and Craig 2005).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human Tonsil
References	<ol style="list-style-type: none"> 1. Dransfield, I. <i>et al.</i> (1992) Interaction of leukocyte integrins with ligand is necessary but not sufficient for function. J Cell Biol. 116:1527-35. 2. Berendt, A. <i>et al.</i> (1992) The binding site on ICAM-1 for plasmodium falciparum-infected erythrocytes overlaps, but is distinct from, the LFA-1- binding site. Cell. 68: 71-81. 3. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. J Pharmacol Exp Ther 321: 656-662. 4. Baratin, M. <i>et al.</i> (2007) Dissection of the role of PfEMP1 and ICAM-1 in the sensing of <i>Plasmodium-falciparum</i>-infected erythrocytes by natural killer cells. PLoS One 2: e228. 5. van Buul, J.D. <i>et al.</i> (2010) Inside-out regulation of ICAM-1 dynamics in TNF-alpha-activated endothelium. PLoS One 5: e11336. 6. Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. J Cell Physiol. 214: 75-83. 7. Di Lorenzo, A. <i>et al.</i> (2011) Endothelial reticulon-4B (Nogo-B) regulates ICAM-1-mediated leukocyte transmigration and acute inflammation. Blood. 117: 2284-95. 8. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. FASEB J. 23: 492-502. 9. Corvaisier, M. <i>et al.</i> (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. J Immunol. 175: 5481-8. 10. Horrocks, P. <i>et al.</i> (2005) PfEMP1 expression is reduced on the surface of knobless Plasmodium falciparum infected erythrocytes. J Cell Sci. 118: 2507-18. 11. Lozanoska-Ochser, B. <i>et al.</i> (2008) Expression of CD86 on human islet endothelial cells facilitates T cell adhesion and migration. J Immunol. 181: 6109-16. 12. Norling, L.V. <i>et al.</i> (2008) Inhibitory control of endothelial galectin-1 on in vitro and in vivo lymphocyte trafficking. FASEB J. 22: 682-90.

13. Baumer, Y. *et al.* (2011) Telomerase-based immortalization modifies the angiogenic/inflammatory responses of human coronary artery endothelial cells. [Exp Biol Med \(Maywood\). 236: 692-700.](#)
14. Lask, A. *et al.* (2011) TCR-independent killing of B cell malignancies by anti-third-party CTLs: the critical role of MHC-CD8 engagement. [J Immunol. 187 \(4\): 2006-14.](#)
15. Sommaggio, R. *et al.* (2012) Multiple Receptors Trigger Human NK Cell-Mediated Cytotoxicity against Porcine Chondrocytes. [J Immunol. 188: 2075-83.](#)
16. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
17. Sumagin R *et al.* (2014) Transmigrated neutrophils in the intestinal lumen engage ICAM-1 to regulate the epithelial barrier and neutrophil recruitment. [Mucosal Immunol. 7 \(4\): 905-15.](#)
18. Sugden SM *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected CD4⁺ T Cells by NK Cells. [J Virol. 91 \(8\): pii: e02442-16.](#)
19. Lennartz, F. *et al.* (2015) Mapping the Binding Site of a Cross-Reactive Plasmodium falciparum PfEMP1 Monoclonal Antibody Inhibitory of ICAM-1 Binding. [J Immunol. 195 \(7\): 3273-83.](#)

Storage Store at +4°C or at -20°C if preferred.

 This product should be stored undiluted.

 Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
- Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#), [DyLight@649](#), [DyLight@680](#), [DyLight@800](#), [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
- Goat Anti Mouse IgG (STAR77...) [HRP](#)
- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
- Human Anti Mouse IgG1 (HCA036...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com
'M316530:180517'

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 25 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)