

Datasheet: MCA1615A647T

Description:	MOUSE ANTI HUMAN CD54:Alexa Fluor® 647
Specificity:	CD54
Other names:	ICAM-1
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	15.2
Isotype:	IgG1
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Pig N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	Human monocytes		

**External Database
Links**

UniProt:

[P05362](#)

[Related reagents](#)

Entrez Gene:

[3383](#)

ICAM1

[Related reagents](#)

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/O-Ag14 myeloma cell line

Specificity

Mouse anti Human CD54 antibody, clone 15.2 recognizes the human CD54 cell surface antigen also known as intracellular Adhesion Molecule -1 (ICAM-1) or Major group rhinovirus receptor.

CD54 is expressed by many cells following activation by inflammatory mediators. It is a 505 amino acid with an additional 27 amino acid signal peptide ~90 kDa single pass type I transmembrane glycoprotein bearing 5 Ig-like C2-type domains domains.

Mouse anti Human CD54 antibody, clone 15.2 blocks CD54 function ([Berendt et al. 1992](#)). Mouse anti Human CD54 antibody, clone 15.2 binds to an epitope on the N-terminal Ig-like domain within a region designated the L43 loop ([Chakravorty and Craig 2005](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Dransfield, I. *et al.* (1992) Interaction of leukocyte integrins with ligand is necessary but not sufficient for function. [J Cell Biol. 116:1527-35.](#)
2. Berendt, A. *et al.* (1992) The binding site on ICAM-1 for plasmodium falciparum-infected erythrocytes overlaps, but is distinct from, the LFA-1- binding site. [Cell. 68: 71-81.](#)
3. Urquhart, P. *et al.* (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther 321: 656-662.](#)
4. Baratin, M. *et al.* (2007) Dissection of the role of PfEMP1 and ICAM-1 in the sensing of *Plasmodium-falciparum*-infected erythrocytes by natural killer cells. [PLoS One 2: e228.](#)
5. van Buul, J.D. *et al.* (2010) Inside-out regulation of ICAM-1 dynamics in TNF-alpha-activated endothelium. [PLoS One 5: e11336.](#)
6. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. [J Cell Physiol. 214: 75-83.](#)
7. Di Lorenzo, A. *et al.* (2011) Endothelial reticulon-4B (Nogo-B) regulates ICAM-1-mediated leukocyte transmigration and acute inflammation. [Blood. 117: 2284-95.](#)
8. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. [FASEB J. 23: 492-502.](#)
9. Corvaisier, M. *et al.* (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. [J Immunol. 175: 5481-8.](#)
10. Horrocks, P. *et al.* (2005) PfEMP1 expression is reduced on the surface of knobless Plasmodium falciparum infected erythrocytes. [J Cell Sci. 118: 2507-18.](#)
11. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells facilitates T cell adhesion and migration. [J Immunol. 181: 6109-16.](#)
12. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on in vitro and in vivo lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
13. Baumer, Y. *et al.* (2011) Telomerase-based immortalization modifies the angiogenic/inflammatory responses of human coronary artery endothelial cells. [Exp Biol Med \(Maywood\). 236: 692-700.](#)
14. Lask, A. *et al.* (2011) TCR-independent killing of B cell malignancies by anti-third-party CTLs: the critical role of MHC-CD8 engagement. [J Immunol. 187 \(4\): 2006-14.](#)

15. Sommaggio, R. *et al.* (2012) Multiple Receptors Trigger Human NK Cell-Mediated Cytotoxicity against Porcine Chondrocytes. [J Immunol. 188: 2075-83.](#)
16. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
17. Sumagin R *et al.* (2014) Transmigrated neutrophils in the intestinal lumen engage ICAM-1 to regulate the epithelial barrier and neutrophil recruitment. [Mucosal Immunol. 7 \(4\): 905-15.](#)
18. Sugden SM *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected CD4⁺ T Cells by NK Cells. [J Virol. 91 \(8\): pii: e02442-16.](#)
19. Lennartz, F. *et al.* (2015) Mapping the Binding Site of a Cross-Reactive Plasmodium falciparum PfEMP1 Monoclonal Antibody Inhibitory of ICAM-1 Binding. [J Immunol. 195 \(7\): 3273-83.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	--	------------------	--	---------------	--

*M300426:170105'

