

Datasheet: MCA1614PE

Description:	MOUSE ANTI HUMAN CD55:RPE
Specificity:	CD55
Other names:	DAF
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	67
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/2

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	K562 cells		
External Database Links	UniProt: P08174 Related reagents		

Entrez Gene:[1604](#) CD55 [Related reagents](#)**Synonyms**

CR, DAF

Specificity

Mouse anti Human CD55 antibody, clone 67 recognizes the human CD55 cell surface antigen, a GPI linked molecule also known as decay accelerating factor (DAF). CD55 is expressed by a wide range of cell types.

CD55 is the complement regulatory protein, decay accelerating factor (DAF) ([Lublin and Atkinson 1989](#)). Human CD55 is a ~70 kDa glycoprotein (in erythrocytes) anchored in the membrane by glycosylphosphatidylinositol tail. In other cells the apparent molecular weight is somewhat larger. It has a substantial content of O-glycans, and also on N-glycan. DAF binds to activated C4b or C3b complement fragments on the cell surface, preventing the assembly and accelerating the decay of both classical and alternative pathways. DAF carries the [Cromer related blood group antigens](#).

DAF has a wide distribution on cells in non-hemopoietic tissues, particularly epithelium and is specifically found at the fetal-maternal interface in placenta ([Holmes et al. 1990](#) and [Yang et al. 2009](#)). Soluble forms of DAF are found, for example, in plasma, saliva and urine ([Medof et al. 1987](#)). The antigen on erythrocytes is pronase and chymotrypsin sensitive, but resistant to trypsin.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Lublin, D.M. & Atkinson, J.P. (1989) Decay-accelerating factor: biochemistry, molecular biology, and function. [Annu Rev Immunol. 7: 35-58.](#)
2. Daniels, G. (1989) Cromer-related antigens--blood group determinants on decay-accelerating factor. [Vox Sang. 56 \(4\): 205-11.](#)
3. Holmes, C.H. et al. (1990) Preferential expression of the complement regulatory protein decay accelerating factor at the fetomaternal interface during human pregnancy. [J Immunol. 144 \(8\): 3099-105.](#)
4. Yang, P. et al. (2009) Expression and modulation of RPE cell membrane complement regulatory proteins. [Invest Ophthalmol Vis Sci. 50: 3473-81.](#)
5. van de Sande, M.G. et al. (2011) Different stages of rheumatoid arthritis: features of the synovium in the preclinical phase. [Ann Rheum Dis. 70: 772-7.](#)
6. Mo, B. et al. (2006) ECC-1 cells: a well-differentiated steroid-responsive endometrial cell line with characteristics of luminal epithelium. [Biol Reprod. 75: 387-94.](#)
7. Araten, D.J. et al. (2005) A quantitative measurement of the human somatic mutation rate. [Cancer Res. 65: 8111-7.](#)
8. de Launay, D. et al. (2010) Silencing the expression of Ras family GTPase homologues decreases inflammation and joint destruction in experimental arthritis. [Am J Pathol. 177: 3010-24.](#)
9. Gheorghe, K.R. et al. (2011) Prostaglandin E2 synthesizing enzymes in rheumatoid arthritis B cells and the effects of B cell depleting therapy on enzyme expression. [PLoS One. ;6: e16378.](#)
10. Kraan, M.C. et al. (2004) T cells, fibroblast-like synoviocytes, and granzyme B+ cytotoxic cells are associated with joint damage in patients with recent onset rheumatoid arthritis. [Ann Rheum Dis. 63: 483-8.](#)
11. van Holten, J. et al. (2005) A multicentre, randomised, double blind, placebo controlled phase II study of subcutaneous interferon beta-1a in the treatment of patients with active rheumatoid arthritis. [Ann Rheum Dis. 64 \(1\): 64-9.](#)
12. Abreu, J.R. et al. (2009) The Ras guanine nucleotide exchange factor RasGRF1 promotes matrix metalloproteinase-3 production in rheumatoid arthritis synovial tissue. [Arthritis Res Ther. 11\(4\):R121.](#)
13. Thurlings, R.M. et al. (2008) Synovial tissue response to rituximab: mechanism of action and

identification of biomarkers of response. [Ann Rheum Dis. 67 \(7\): 917-25.](#)

14. Vos, K. *et al.* (2007) Early effects of rituximab on the synovial cell infiltrate in patients with rheumatoid arthritis. [Arthritis Rheum. 56 \(3\): 772-8.](#)

15. Edginton S *et al.* (2016) Effects of Rituximab and Infliximab Treatment on Carboxypeptidase B and Its Substrates in RA Synovium. [J Rheumatol. 43 \(5\): 846-54.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

12 months from date of reconstitution.

Health And Safety Information

Material Safety Datasheet documentation #10075 available at:
10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South

Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M312129:180126'

Printed on 01 Aug 2018