

Datasheet: MCA1757T

Description:	MOUSE ANTI HUMAN MYELOPEROXIDASE
Specificity:	MYELOPEROXIDASE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2C7
Isotype:	IgG1
Quantity:	20 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/1000 - 1/5000
Immunohistology - Paraffin	▪			1/500 - 1/1000
ELISA	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) **Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

Target Species	Human
Species Cross Reactivity	Reacts with: Dog Does not react with:Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	Human myeloperoxidase.
External Database Links	<p>UniProt: P05164 Related reagents</p> <p>Entrez Gene: 4353 MPO Related reagents</p>
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse X63 AG8-653 myeloma cell line.
Specificity	<p>Mouse anti Human myeloperoxidase antibody, clone 2C7 recognizes human myeloperoxidase (MPO). MPO is an important component of azurophilic granules in neutrophils, being involved in microbicidal processes. The protein is a multimer of 2 heavy chains (55 kDa) and two light chains (15 kDa), the heavy chains being linked by a disulphide bond.</p> <p>Mouse anti Human Myeloperoxidase antibody, clone 2C7 recognizes native MPO in Western blots, and the heavy chain following boiling of the sample. Mouse anti Human Myeloperoxidase antibody, clone 2C7 also recognizes recombinant MPO in western blots and weakly in ELISA.</p> <p>Mouse anti Human myeloperoxidase antibody, clone 2C7 may be of value in the study of myeloid cells and myeloid leukaemias by flow cytometry following cell permeabilization. Mouse anti Human myeloperoxidase antibody, clone 2C7 did not recognize rat MPO by ELISA (Patry et al. 2003).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Bone marrow
References	<ol style="list-style-type: none"> Audrain, M.A. <i>et al.</i> (1997) Anti-native and recombinant myeloperoxidase monoclonals and human autoantibodies. Clin Exp Immunol. 107 (1): 127-34. Patry, Y.C. <i>et al.</i> (2003) Difference in antigenic determinant profiles between human and rat myeloperoxidase. Clin Exp Immunol. 132 (3): 505-8. Zhang, N. <i>et al.</i> (2008) Different types of T-effector cells orchestrate mucosal inflammation in chronic sinus disease. J Allergy Clin Immunol. 122: 961-8. Sloane, A.J. <i>et al.</i> (2005) Proteomic analysis of sputum from adults and children with cystic fibrosis and from control subjects. Am J Respir Crit Care Med. 172: 1416-26. Luo, B. <i>et al.</i> (2013) Immunopathology features of chronic rhinosinusitis in high-altitude dwelling Tibetans. Allergy Rhinol (Providence). 4: e69-76. Behnen, M. <i>et al.</i> (2014) Immobilized immune complexes induce neutrophil extracellular trap release by human neutrophil granulocytes via FcγRIIIB and Mac-1. J Immunol. 193 (4): 1954-65. Villiers, E. <i>et al.</i> (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. Vet Clin Pathol. 35 (1): 55-71. Gelain, M.E. <i>et al.</i> (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. Vet Immunol Immunopathol. 159 (1-2): 91-6. Wang, H. <i>et al.</i> (2016) Circulating Level of Neutrophil Extracellular Traps Is Not a Useful Biomarker for Assessing Disease Activity in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. PLoS One. 11 (2): e0148197. Rai, A.K. <i>et al.</i> (2017) Exonal switch down-regulates the expression of CD5 on blasts of acute T cell leukaemia. Clin Exp Immunol. 190 (3): 340-350. Novacco, M. <i>et al.</i> (2016) Prognostic factors in canine acute leukaemias: a retrospective study.

[Vet Comp Oncol. 14 \(4\): 409-16.](#)

12. Villiers, E. *et al.* (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. [Vet Clin Pathol. 35 \(1\): 55-71.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Human Anti Mouse IgG1 (HCA036...) [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#),
[DyLight@649](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M317102:180619'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)