

Datasheet: MCA1701F

Description:	MOUSE ANTI p53 (aa20-25):FITC
Specificity:	p53 (aa20-25)
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	DO-1
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/2 - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

Target Species	Human								
Species Cross Reactivity	Reacts with: Bovine, Cat, Horse, Green Monkey Does not react with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.								
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid								
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525		
Fluorophore	Excitation Max (nm)	Emission Max (nm)							
FITC	490	525							
Preparation	Purified IgG prepared by affinity chromatography on Protein G								
Buffer Solution	Phosphate buffered saline								
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin								
Approx. Protein Concentrations	IgG concentration 0.1mg/ml								
Immunogen	Recombinant human p53.								

External Database Links	<p>UniProt: P04637 Related reagents</p> <p>Entrez Gene: 7157 TP53 Related reagents</p>
Synonyms	P53
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Human p53 antibody, clone DO-1 recognizes the human p53 tumor suppressor protein, also known as cellular tumor antigen p53 or NY-CO-13. Clone DO-1 binds to both wild type and mutant forms of the p53 protein found in various malignancies (Kern et al. 1992). p53 is important in multicellular organisms, where it regulates cell cycle progression to allow DNA repair or apoptosis in the case of irreparably damaged cells (Haupt et al. 2003) and thus functions as a tumor suppressor that is involved in preventing cancer. Mutations in the p53 gene are found in about half the cases of human cancer (Joerger andFersht 2007)</p> <p>Mouse anti Human p53 antibody, clone DO-1 recognizes an epitope at the N-terminal end of p53 between amino acids 20-25,common to isoforms 1-3 of p53.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Vojtěšek B <i>et al.</i> (1992) An immunochemical analysis of the human nuclear phosphoprotein p53. New monoclonal antibodies and epitope mapping using recombinant p53. J Immunol Methods. 151 (1-2): 237-44. 2. Sironi, G. <i>et al.</i> (1999) p53 protein expression in conjunctival squamous cell carcinomas of domestic animals. Vet Ophthalmol. 2 (4): 227-231. 3. Levesque, M.A. <i>et al.</i> (1995) Time-resolved immunofluorometric assay of p53 protein. Clin Chem. 41 (12 Pt 1): 1720-9. 4. Bonsing, B.A. <i>et al.</i> (1997) Specificity of seven monoclonal antibodies against p53 evaluated with Western blotting, immunohistochemistry, confocal laser scanning microscopy, and flow cytometry. Cytometry. 28 (1): 11-24. 5. Carvalho, T. <i>et al.</i> (2009) Immunohistochemical evaluation of vascular urinary bladder tumors from cows with enzootic hematuria. Vet Pathol. 46 (2): 211-21. 6. Phillips, A. <i>et al.</i> (2010) HDMX-L is expressed from a functional p53-responsive promoter in the first intron of the HDMX gene and participates in an autoregulatory feedback loop to control p53 activity. J Biol Chem. 285 (38): 29111-27. 7. Bergman, L.M. <i>et al.</i> (2009) CtBPs promote cell survival through the maintenance of mitotic fidelity. Mol Cell Biol. 29: 4539-51. 8. Hietanen, S. <i>et al.</i> (2000) Activation of p53 in cervical carcinoma cells by small molecules. Proc Natl Acad Sci U S A. 97 (15): 8501-6. 9. Phelps, M. <i>et al.</i> (2003) p53-independent activation of the hdm2-P2 promoter through multiple transcription factor response elements results in elevated hdm2 expression in estrogen receptor alpha-positive breast cancer cells. Cancer Res. 63: 2616-23.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p>

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:FITC \(MCA929F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M302097:170109'

Printed on 01 Aug 2018