

Datasheet: MCA1768XZ

Description:	RAT ANTI MOUSE CD8:Preservative Free
Specificity:	CD8
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	YTS169.4
Isotype:	IgG2b
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: P01731 Related reagents

[P10300](#) [Related reagents](#)

Entrez Gene:

[12525](#) Cd8a [Related reagents](#)

[12526](#) Cd8b1 [Related reagents](#)

Synonyms Cd8b1, Ly-3, Lyt2, Lyt-2, Lyt3, Lyt-3

Specificity **Rat anti Mouse CD8 antibody, clone YTS169.4** recognizes the murine CD8 cell surface antigen, expressed by a subset of T lymphocytes.

Rat anti Mouse CD8 antibody, clone YTS169.4 exhibits depleting activity when used *in vivo*.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Cobbold, S.P. *et al.* (1990) The induction of skin graft tolerance in major histocompatibility complex-mismatched or primed recipients: primed T cells can be tolerized in the periphery with anti-CD4 and anti-CD8 antibodies. [Eur J Immunol. 20 \(12\): 2747-55.](#)
2. Bemelman, F. *et al.* (1998) Bone marrow transplantation induces either clonal deletion or infectious tolerance depending on the dose. [J Immunol. 160 \(6\): 2645-8.](#)
3. Cobbold SP *et al.* (1984) Therapy with monoclonal antibodies by elimination of T-cell subsets *in vivo*. [Nature. 312 \(5994\): 548-51.](#)
4. Wise, M.P. *et al.* (1998) Linked suppression of skin graft rejection can operate through indirect recognition. [J Immunol. 161 \(11\): 5813-6.](#)
5. Higgins, L.M. *et al.* (1999) Regulation of T cell activation in vitro and in vivo by targeting the OX40-OX40 ligand interaction: amelioration of ongoing inflammatory bowel disease with an OX40-IgG fusion protein, but not with an OX40 ligand-IgG fusion protein. [J Immunol. 162 \(1\): 486-93.](#)
6. Scotland, R.S. *et al.* (2011) Sex-differences in resident immune cell phenotype underlies more efficient acute inflammatory responses in female mice. [Blood. 118: 5918-27.](#)
7. Matsubara, K. *et al.* (2016) Immune activation during the implantation phase causes preeclampsia-like symptoms via the CD40-CD40 ligand pathway in pregnant mice. [Hypertens Res. 39 \(6\): 407-14.](#)
8. Jaffar, Z. *et al.* (2002) A key role for prostaglandin I2 in limiting lung mucosal Th2, but not Th1, responses to inhaled allergen. [J Immunol. 169 \(10\): 5997-6004.](#)
9. Zirger, J.M. *et al.* (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFN γ , perforin, and TNF α , and due to the elimination of transduced cells. [Mol Ther. 20 \(4\): 808-19.](#)
10. Abd-elhakim, Y.M. *et al.* (2016) Hemato-immunologic impact of subchronic exposure to melamine and/or formaldehyde in mice. [J Immunotoxicol. 13 \(5\): 713-22.](#)
11. Ismail, S.A.A. (2017) Ameliorative Potential of *Spirulina platensis* against Lead Acetate Induced Immuno-Suppression and Kidney Apoptosis in Rats [Ann Clin Pathol 5\(5\): 1120.](#)

Storage Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10162 available at:
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M315394:180503'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)