

Datasheet: MCA1846PE

Description:	HAMSTER ANTI MOUSE CD81:RPE
Specificity:	CD81
Other names:	TAPA-1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	Eat2
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	38C13, murine B cell line.		

**External Database
Links**

UniProt:

[P35762](#) [Related reagents](#)

Entrez Gene:

[12520](#) Cd81 [Related reagents](#)

Synonyms

Tapa1

Fusion Partners

Spleen cells from immunised Armenian hamsters were fused with cells of the mouse PX3-Ag.8.653 myeloma cell line.

Specificity

Hamster anti Mouse CD81 antibody, clone Eat2 recognizes mouse and rat CD81, also known as TAPA-1 or Target of the antiproliferative antibody 1. CD81 is a 236 amino acid ~26 kDa multipass transmembrane protein belonging to the TM4SF family ([UniProt: P35762](#)). In rodents CD81 is expressed at much higher levels on resting B cells than on T cells, although increased expression on T cells is found following activation. Hamster anti Mouse CD81 antibody, clone Eat2 induces homotypic aggregation of B cells and inhibits anti Ig and IL-4 induced proliferation ([Maecker *et al.* 2000](#)). Eat 2 requires the presence of both extracellular loops of TAPA-1 for binding.

Mice lacking CD81 demonstrate reduced fertility through impaired oocyte-sperm fusion, double knockout CD81^{-/-} CD9^{-/-} mice are completely infertile suggesting complimentary roles in oocyte-sperm fusion ([Rubenstein *et al.* 2006](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

- Clark, K.L. *et al.* (2001) PGRL is a major CD81-associated protein on lymphocytes and distinguishes a new family of cell surface proteins. [J Immunol. 167 \(9\): 5115-21.](#)
- Maecker, H.T. *et al.* (2000) Differential expression of murine CD81 highlighted by new anti-mouse CD81 monoclonal antibodies. [Hybridoma 19: 15-22.](#)
- Conde-Vancells, J. *et al.* (2010) Candidate biomarkers in exosome-like vesicles purified from rat and mouse urine samples. [Proteomics Clin Appl. 4 \(4\): 416-25.](#)
- Conde-Vancells, J. *et al.* (2008) Characterization and comprehensive proteome profiling of exosomes secreted by hepatocytes. [J Proteome Res. 7: 5157-66.](#)
- Takeda, Y. *et al.* (2008) Double deficiency of tetraspanins CD9 and CD81 alters cell motility and protease production of macrophages and causes chronic obstructive pulmonary disease-like phenotype in mice. [J Biol Chem. 283: 26089-97.](#)
- Suzuki, M. *et al.* (2009) Tetraspanin CD9 negatively regulates lipopolysaccharide-induced macrophage activation and lung inflammation. [J Immunol. 182: 6485-93.](#)
- Ha, C.T. *et al.* (2005) Binding of pregnancy-specific glycoprotein 17 to CD9 on macrophages induces secretion of IL-10, IL-6, PGE2, and TGF-beta1. [J Leukoc Biol. 77: 948-57.](#)
- Pan, Q. *et al.* (2011) Hepatic cell-to-cell transmission of small silencing RNA can extend the therapeutic reach of RNA interference (RNAi). [Gut. 61: 1330-9.](#)
- Jin, Y. *et al.* (2013) Statins decrease lung inflammation in mice by upregulating tetraspanin CD9 in macrophages. [PLoS One. 8: e73706.](#)
- Royo, F. *et al.* (2013) Transcriptome of extracellular vesicles released by hepatocytes. [PLoS One. 8: e68693.](#)
- Owens, D.M. and Watt, F.M. (2001) Influence of beta1 integrins on epidermal squamous cell carcinoma formation in a transgenic mouse model: alpha3beta1, but not alpha2beta1, suppresses malignant conversion. [Cancer Res. 61: 5248-54.](#)

12. Jin, Y. *et al.* (2018) Double deletion of tetraspanins CD9 and CD81 in mice leads to a syndrome resembling accelerated aging. [Sci Rep. 8 \(1\): 5145.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	--

Shelf Life	12 months from date of reconstitution.
-------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M301016:170106'

Printed on 20 Jun 2018