

Datasheet: MCA1940F

Description:	MOUSE ANTI HUMAN CD19:FITC
Specificity:	CD19
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	LT19
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
External Database Links	UniProt: P15391Related reagents Entrez Gene: 930 CD19 Related reagents		

Specificity	Mouse anti Human CD19 antibody, clone LT19 recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells (de Rie et al. 1989). CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 (Bradbury et al. 1992), CD9, CD81 and CD82 (Horváth et al. 1998). CD19 is implicated in the down-regulation of B cell growth and proliferation (Pezzutto et al. 1987).
Flow Cytometry	Use 10ul of the suggested working dilution to label 100ul of lysed whole peripheral blood.
References	- Hughes, G.J. <i>et al.</i> (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 <i>in vivo</i>. AIDS. 21: 1507-13. - Allen, J.S. <i>et al.</i> (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. Diabetes. 58: 138-45. - McIntosh, K. <i>et al.</i> (2006) The immunogenicity of human adipose-derived cells: temporal changes <i>in vitro</i>. Stem Cells. 24: 1246-53. - Sengstake, S. <i>et al.</i> (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. Int Immunol. 18 (7): 1171-8. - Villarroel Dorrego, M. <i>et al.</i> (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. Br J Dermatol. 154: 231-8. - Franz, B. <i>et al.</i> (2011) <i>Ex vivo</i> characterization and isolation of rare memory B cells with antigen tetramers. Blood. 118: 348-57. - Lacal, P.M. <i>et al.</i> (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. J Pharmacol Exp Ther. 347: 164-72. - Franz, B. <i>et al.</i> (2011) <i>Ex vivo</i> characterization and isolation of rare memory B cells with antigen tetramers. Blood. 118: 348-57. - Girbl, T. <i>et al.</i> (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. Cancer Res. 73: 561-70. - Hertzberg, L. <i>et al.</i> (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. Blood. 115: 1006-17. - Kakko, T. <i>et al.</i> (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. Diab Vasc Dis Res. 8: 221-8. - Dorviguit, D. <i>et al.</i> (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. MAbs. 4 (4): 488-96. - Karlsen, M. <i>et al.</i> (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. Scand J Immunol. 82 (6): 523-31.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product

contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301381:170109'

Printed on 02 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)